



Vegetation Classification and Mapping of Tallgrass Prairie National Preserve

Project Report

Natural Resource Report NRR/HTLN/NRR—2011/346



ON THE COVER

Tallgrass prairie landscape at the Tallgrass Prairie National Preserve; view of stone school with wild rye (*Elymus canadensis*), wild alfalfa (*Psoraleum tenuifolium*), and blacksamson echinacea (*Echinacea angustifolia*) seedheads in the foreground.

Photograph by: Kelly Kindscher

Vegetation Classification and Mapping of Tallgrass Prairie National Preserve

Project Report

Natural Resource Report NRR/HTLN/NRR—2011/346

Kelly Kindscher^{1*}
Hayley Kilroy¹
Jennifer Delisle¹
Quinn Long¹
Hillary Loring¹
Kevin Dobbs²
Jim Drake³

¹Kansas Natural Heritage Inventory
Kansas Biological Survey
University of Kansas
2101 Constant Ave.
Lawrence, KS 66047

²Kansas Applied Remote Sensing Program
Kansas Biological Survey
University of Kansas
2101 Constant Ave.
Lawrence, KS 66047

³NatureServe
P.O. Box 9354
St. Paul, MN 55109

*contact kindscher@ku.edu

April 2011

U.S. Department of the Interior
National Park Service
Natural Resource Program Center
Fort Collins, Colorado

The National Park Service, Natural Resource Program Center publishes a range of reports that address natural resource topics of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Report Series is used to disseminate high-priority, current natural resource management information with managerial application. The series targets a general, diverse audience, and may contain NPS policy considerations or address sensitive issues of management applicability.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data. Data in this report were collected and analyzed using methods based on established, peer-reviewed protocols and were analyzed and interpreted within the guidelines of the protocols.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from <http://web.ku.edu/~kindscher/tallgrass> and the Natural Resource Publications Management website (<http://www.nature.nps.gov/publications/nrpm/>).

Please cite this publication as:

Kindscher, K., H. Kilroy, J. Delisle, Q. Long, H. Loring, K. Dobbs, and J. Drake. 2011. Vegetation mapping and classification of Tallgrass Prairie National Preserve: Project report. Natural Resource Report NRR/HTLN/NRR—2011/346. National Park Service, Fort Collins, Colorado.



Contents

	Page
Figures.....	v
Tables.....	viii
Appendices.....	xi
Executive Summary.....	xiii
Introduction.....	1
Tallgrass Prairie National Preserve Vegetation Mapping Project.....	1
USGS-NPS Vegetation Mapping Program.....	1
<i>Vegetation Mapping Program Standards</i>	2
<i>Tallgrass Prairie National Preserve</i>	2
Project Statistics.....	4
Methods.....	5
Planning, Data Gathering and Coordination.....	5
<i>Field Survey</i>	5
<i>Vegetation Classification</i>	7
Digital Imagery and Interpretation.....	9
<i>Accuracy Assessment</i>	10
Results.....	15
Vegetation Classification.....	15
Digital Imagery and Interpretation.....	15
<i>Vegetation Map</i>	15
<i>Accuracy Assessment</i>	15
Vegetation Associations.....	19
<i>Mapped Unit Name: Bur Oak Woodland</i>	19

Contents (continued)

	Page
<i>Mapped Unit Name: Successional Forest</i>	22
<i>Mapped Unit Name: Native Tallgrass Prairie</i>	24
<i>Mapped Unit Name: Rocky Mixed Prairie</i>	26
<i>Mapped Unit Name: Wet Ravine Vegetation</i>	29
<i>Mapped Unit Name: Wet Ravine Vegetation - Seeps and Springs</i>	32
<i>Mapped Unit Name: Smooth Brome</i>	34
<i>Common Name: Restored Prairie</i>	36
<i>Common Name: Weedy</i>	38
Discussion.....	41
Field Survey.....	41
NVC Classification.....	41
Digital Imagery and Interpretation	41
Accuracy Assessment	41
Future Recommendations	41
Research Opportunities.....	42
Literature Cited.....	43

Figures

	Page
Figure 1. Location of Tallgrass Prairie National Preserve in Chase County, Kansas.....	3
Figure 2. Map of the vegetation project boundary and park boundary.....	6
Figure 3. Locations of all vegetation plots collected at TAPR in 2008.	8
Figure 4. Locations of all accuracy assessment points collected at TAPR in 2009.....	13
Figure 5. Vegetation map of Tallgrass Prairie National Preserve and environs.....	17
Figure 6. Bur Oak Woodland at Tallgrass Prairie National Preserve, July 2009.	19
Figure 7. Successional Forest at Tallgrass Prairie National Preserve, July 2009.	22
Figure 8. Native Tallgrass Prairie at Tallgrass Prairie National Preserve, July 2009.....	24
Figure 9. Rocky Mixed Prairie at Tallgrass Prairie National Preserve, July 2009.	26
Figure 10. Wet Ravine at Tallgrass Prairie National Preserve, July 2009.....	29
Figure 11. Seep at Tallgrass Prairie National Preserve, July 2009. Darker green in midground is <i>Eleocharis spp.</i> , tall seedheads are <i>Scirpus pendulus</i> , and dark brown seedheads in foreground and background are <i>Rumex spp.</i>	32
Figure 12. Restored Prairie at Tallgrass Prairie National Preserve, July 2008.....	36

Tables

	Page
Table 1. Target number of AA samples per map class based on number of polygons and area.	11
Table 2. Summary of the AA statistics used at TAPR.	11
Table 3. Map units identified at TAPR, with their total frequency and acreage.	16
Table 4. Overstory Tree Species (DBH>15 cm) within Bur Oak Woodland. (Data collected by Heartland Inventory and Monitoring Program, n=10 plots).....	20
Table 5. Overstory Tree Species (DBH>15 cm) within Successional Forest. (Data collected by the Heartland Network breeding bird monitoring (n=4 plots).	23
Table 6. Average percent cover of the top twenty most common species in plots within Native Tallgrass Prairie. (Data collected by the Heartland Network in 2008, n=23 plots.)	25
Table 7. Average percent cover of the top twenty most common species in plots within Rocky Mixed Prairie. (Data collected by KBS in 2008, n=10 plots.)	27
Table 8. Average percent cover of the top twenty most common species in plots within Wet Ravines. (Data collected by KBS in 2008, n=10 plots; note many areas have woody encroachment)	30
Table 9. Average percent cover of the top twenty most common species in plots within Seeps and Springs. (Data collected by KBS in 2008, n=10 plots.)	33
Table 10. Average percent cover of the top twenty most common species in plots within Smooth Brome Haymeadows. Note that all species other than <i>Bromus inermis</i> are under 3% cover. (Data collected by KBS in 2008, n=10 plots.).....	35
Table 11. Average percent cover of the top twenty most common species in plots within Restored Prairies. (Data collected by KBS in 2008, n=19 plots.)	37
Table 12. Average percent cover of the top nineteen most common species in plots within Weedy Areas. (Data collected by KBS in 2008, n=4 plots.)	39

Appendices

	Page
Appendix A: contingency Table for Vegetation Mapping at TAPR	47
Appendix B: Example of a Plot Sampling Form	49
Appendix C: Tallgrass Prairie National Preserve Dichotomous Key to Vegetation Associations	51
Appendix D: Example of an Accuracy Assessment Sampling Form	53
Appendix E: Tallgrass Prairie National Preserve Species List	54

Executive Summary

The Tallgrass Prairie National Preserve (TAPR) encompasses 10,894 acres in eastern Kansas, just north of Strong City. This park unit was created on November 12, 1996 and is the first to protect a nationally significant example of the once vast tallgrass prairie ecosystem. Of the 400,000 square miles of tallgrass prairie that once covered the North American continent, less than four percent remains, primarily in the Flint Hills. The park unit is primarily rocky upland prairies and deep-soiled prairies in the lowlands. It also contains some wet prairie ravines, riparian forests and some former cropland and restored prairie.

A three-year program was initiated to complete the task of mapping and classifying the vegetation at TAPR. The Kansas Biological Survey (KBS) in conjunction with NatureServe developed a vegetation classification using the National Vegetation Classification System and produced a digital vegetation map. To classify the vegetation, plots located throughout TAPR were sampled during the summer of 2008. Additional data were obtained from vegetation plots sampled by the Inventory & Monitoring program in 2006. Analysis of the plot data by KBS produced 12 map units (eight vegetated and four land-use) which are directly matched to corresponding plant associations and land-use classes. Descriptions and a field key for all plant communities of TAPR are included in this report. Draft maps were printed, field tested, reviewed and revised. Accuracy assessment (AA) data points were collected on 112 data points in 2009 by KBS and used to verify the map's accuracy.

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

Introduction

Tallgrass Prairie National Preserve Vegetation Mapping Project

The Tallgrass Prairie National Preserve (TAPR) Vegetation Mapping Project was organized and coordinated by the Kansas Biological Survey (KBS) at the University of Kansas, in cooperation with NatureServe, in accordance with the standards set forth by the U.S. Geological Survey (USGS) – National Park Service (NPS) Vegetation Mapping Program.

The TAPR Vegetation Mapping Project was initiated because the preserve protects a nationally significant example of the once vast tallgrass prairie ecosystem. Of the 400,000 square miles of tallgrass prairie that once covered the North American continent, less than four percent remains, primarily in the Flint Hills. Although the Tallgrass Prairie National Preserve had been mapped at a coarse level (tallgrass prairie vs. riparian forest area), a more accurate map was needed to break out grassland types, rock outcrops, replanted vegetation, water, and other landmarks to National Vegetation Classification alliance. A unified objective classification, such as outlined in the National Park Service's Vegetative Mapping Program, can become a valuable aid to the preserve for the use in vegetation management, grazing, fire, and monitoring wetlands and wildlife. Since the National Park Service is charged with conserving, protecting, and interpreting the resources of this prairie landscape, an accurate and detailed vegetation map and data layers for a GIS is very useful for management purposes.

Common to all Vegetation Mapping Program projects, the three major components of the TAPR Vegetation Mapping Project are vegetation classification, vegetation mapping, and map accuracy assessment. In this report we discuss each of these fundamental components in detail.

USGS-NPS Vegetation Mapping Program

The National Vegetation Mapping Program is an interagency initiative established to inventory, classify, describe, and map vegetation in National Park units and other areas across the United States. It is administered by the USGS Center for Biological Informatics and the NPS Natural Resources Information Division, and provides baseline vegetation information to the NPS Inventory and Monitoring Program (I&M).

Vegetation Mapping Program scientists developed procedures for classification, mapping, and accuracy assessment (The Nature Conservancy [TNC] and Environmental Systems Research Institute [ESRI] 1994a).

Use of the National Vegetation Classification System (NVCS) as the standard vegetation classification system is central to fulfilling the goals of this national program. This system:

- is vegetation based;
- uses a systematic approach to classify a continuum;
- emphasizes natural and existing vegetation;
- uses a combined physiognomic-floristic hierarchy;
- identifies vegetation units based on both qualitative and quantitative data;
- is appropriate for mapping at multiple scales.

The use of the NVCS and the USGS-NPS vegetation mapping protocols facilitates effective resource stewardship by ensuring compatibility and widespread use of the information throughout the NPS as

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

well as by other federal and state agencies. These vegetation maps and associated information support a wide variety of resource assessment, park management, and planning needs. In addition they can be used to provide a structure for framing and answering critical scientific questions about vegetation communities and their relationship to environmental conditions and ecological processes across the landscape.

The NVCS has primarily been developed and implemented by The Nature Conservancy (TNC) and the network of state natural heritage programs over the past twenty years (TNC and ESRI 1994a; Grossman et al. 1998). The NVCS is currently supported and endorsed by multiple federal agencies, the Federal Geographic Data Committee (FGDC), NatureServe, state heritage programs, and the Ecological Society of America. Refinements to the classification occur in the process of application, leading to ongoing proposed revisions that are reviewed both locally and nationally.

Vegetation Mapping Program Standards

The NPS I&M Program established guidance and standards for all vegetation mapping projects in a series of documents:

Protocols

- documenting a National Vegetation Classification System (TNC and ESRI 1994a);
- standards for field methods and mapping procedures (TNC and ESRI 1994b);
- producing rigorous and consistent accuracy assessment procedures (TNC et al. 1994);
- establishing standards for using existing vegetation data (TNC 1996);

Standards

- National Vegetation Classification Standard (FGDC 1997);
- Spatial Data Transfer Standard (FGDC 1998b);
- Content Standard for Digital Geospatial Metadata (FGDC 1998a);
- United States National Map Accuracy Standards (USGS 1999);
- Integrated Taxonomic Information System (<http://www.itis.gov/>);
- program-defined standards for map attribute accuracy and minimum mapping unit.

These documents are available on the USGS-NPS Vegetation Program Web site (<http://biology.usgs.gov/npsveg/standards.html>).

Tallgrass Prairie National Preserve

In November of 1996, federal legislation was passed creating the 10,894 acre Tallgrass Prairie National Preserve in the Flint Hills region of Kansas, located in Chase County, Kansas, outside of Strong City. Ownership of the park is a unique private/public partnership between the National Park Service and The Nature Conservancy. The preserve protects a nationally significant example of the once vast tallgrass prairie ecosystem.

Tallgrass Prairie National Preserve was created on November 12, 1996 to "...preserve, protect, and interpret for the public an example of a tallgrass prairie ecosystem..." and to "...preserve and interpret for the public the historic and cultural values represented on the Spring Hill Ranch."

The preserve is a designated National Historic Landmark that, "...outstandingly represents the transition from the open range to the enclosed holdings of the large cattle companies in the 1880's." In addition, the nomination notes that "...the ranch lands (contributing site) have retained a high




Figure 1. Location of Tallgrass Prairie National Preserve in Chase County, Kansas.

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

level of integrity in all areas of consideration: location, setting, feeling, design, materials, workmanship, as well as association” (Wolfenbarger and Nimz 1996)

The preserve was established to provide visitors access to the cultural and natural features of the Spring Hill/Z Bar Ranch. The preserve actively uses large grazers, i.e. bison and cattle, to manage the natural landscape and represent the cultural landscape.

Project Statistics

Field Work Summers of 2008 and 2009:

Plot Sampling = 77 Plots:

54 plots sampled in July 2008 by the Kansas Biological Survey staff

23 plots sampled during the 2008 growing season by the Heartland Network

Accuracy Assessment Points = 132

112 points to assess vegetation classification

20 additional points to assess presence of seeps

All collected in July 2009 by the Kansas Biological Survey staff

Classification:

6 NVC Plant Associations

2 Park Special Vegetation Classes

4 Non-Vegetated Land-Use Classes

GIS Database 2006 – 2009:

Project Size = 18,805 acres (7610.11 hectares)

Tallgrass Prairie National Preserve = 10,894 acres (4408.65 hectares)

Environs = 7,911 acres (3201.47 hectares)

Base Imagery acquired from the NPS:

Fall 2005 IKONOS image

Ancillary Imagery acquired by the Kansas Applied Remote Sensing Program, a program of the Kansas Biological Survey:

June and September of 2008 IKONOS image

2003, 2004, 2005, 2006, and 2008 United States Department of Agriculture (USDA)

Farm Service Agency National Agriculture Imagery Program (NAIP)

2002 and 1991 United States Geological Survey (USGS) Digital Orthophoto Quarter

Quads (DOQQs)

1938 panchromatic imagery

Minimum Mapping Unit = 0.5 hectare

Total Size = 569 Polygons

Average Polygon Size = 33.0 acres (13.35 hectares)

Overall Thematic Accuracy = 92.0%

Project Completion Date: 06/30/2010

Methods

The entire map extent is 18,730 acres, of which the park comprises 10,861 acres (Figure 2). The national standard minimum mapping unit is 0.5 ha, but some polygons, notably those of seeps or thin-soiled rocky areas, are smaller than the minimum mapping unit.

The vegetation mapping project at Tallgrass Prairie National Preserve was considered to be in the “medium park” category based on the overall size of the project area (TNC 1994b). As such, the standard methodology for sampling and mapping is to visit the entire park and select representative sites. It is assumed that these sites will sufficiently characterize the vegetation types and explain their distribution across the park without having to survey each stand of vegetation. Based on this approach the assignment of responsibilities was divided into five major tasks, including the following:

1. Plan, gather data, and coordinate tasks;
2. Survey TAPR to understand and sample the vegetation;
3. Classify the vegetation using the field data to NVC standard associations and alliances and crosswalk these to recognizable map units;
4. Acquire current digital imagery and interpret the vegetation from these using the classification scheme and a map unit crosswalk;
5. Assess the accuracy of the final map product.

All protocols for this project as outlined in the following sections can be found in documents produced by TNC and ESRI (1994a, 1994b) and TNC et al. (1994) for the USGS-NPS Vegetation Mapping Program. These documents can be found at: <http://biology.usgs.gov/npsveg>.

Planning, Data Gathering and Coordination

A scoping meeting was held in June 2007 with all project participants (Kansas Biological Survey, NatureServe, Tallgrass Prairie National Preserve staff, NPS Heartland Network staff, Homestead National Monument staff, National Park Service National Vegetation Mapping staff, and The Nature Conservancy). KBS was responsible for plot sampling and reconnaissance visits of potential community types of TAPR. KBS was also responsible for entering these data into a digital database, classifying these data, and providing a list and global descriptions for the TAPR plant associations. KBS was responsible for the imagery interpretation and creating a digital vegetation map and spatial database. NatureServe reviewed and evaluated the draft classification and wrote vegetation descriptions for all associations. KBS created a vegetation key, and conducted accuracy assessment of the vegetation map. NatureServe and TAPR staff provided logistical and technical support, and helped coordinate activities.

Field Survey

The field methods used by the Kansas Natural Heritage Inventory in sampling and classifying the vegetation followed the methodology outlined by the USGS-BRD/NPS Vegetation Mapping




Figure 2. Map of the vegetation project boundary and park boundary.

Program and the NVC (Grossman et al. 1994, Grossman et al. 1998). The application of these methods to Tallgrass Prairie National Preserve is outlined below.

Vegetation data were collected in characteristic plots by KBS in July 2008 and from additional plot information acquired from the Heartland Inventory and Monitoring Network (Figure 3). Characteristic plots were located in areas that were visually representative of the preliminary vegetation categories. Plots were 100 m² in area, and GPS coordinates were recorded with a Garmin receiver. To maintain consistency with other projects in Kansas, plots were square. For a couple of the seeps in was actually necessary to make rectangular plots totaling 100m² in area due to irregular shapes of the feature. The accuracy for all of the recorded points ranged from 1-7 meters in horizontal accuracy, as recorded by the GPS receiver. Ten plots were sampled for most preliminary vegetation types: rocky mixed prairie, restored prairie, riparian vegetation, and brome fields. For “weedy” vegetation, only four characteristic plots were sampled due to the low abundance of the vegetation type. Ten plots were also sampled for springs and seeps, which were not characterized as a polygon vegetation type, but were mapped as points from Kansas Geological Survey data.

All plants found within the characteristic plots were identified to species level where possible. In a few cases, identification was only possible to the genus level (i.e., non-reproductive *Muhlenbergia* and *Carex* species). Visual estimates of percent cover were made for all species, including live material and the current year’s standing dead. To maintain consistency with local vegetation surveys and other work of KBS, a continuous range of possible cover estimates was used, rather than cover classes. Plants found to cover at least one half of one percent of the plot were assigned one percent (0.01) and those with less than one half of one percent a “trace” (T). Also to maintain consistency with published accounts and similar projects in the region, species were assigned names following the Flora of the Great Plains (McGregor and Barkley 1986). An updated synonymy was completed when data were entered into the PLOTS database. Noteworthy surrounding vegetation, slopes, unusual soil features, and noticeable use by animals were also noted at each plot. Most plots were on a gentle slope with an A horizon of silty clay loam soils.

Additional plot data were obtained from the Heartland Inventory and Monitoring Network (James et al 2009). Tallgrass prairie sampling was conducted in 2008 on ten permanent plots. Each plot comprises two, 50 m long transects with ten sets of nested subplots systematically arranged. Working from the smallest to the largest plot, all herbaceous, woody shrub and tree seedling and sapling species were identified. Foliar cover was estimated in the 100 m² subplot using a modified Daubenmire (1959) scale. Forest overstory data were obtained from a breeding bird survey conducted in 2008, in which overstory trees were tallied in 18 plots (Peitz et al 2008).

Vegetation Classification

Upon completion of field surveys, all recorded data were entered into the NPS PLOTS database (TNC 1997), a Microsoft Access-derived program. The PLOTS database was developed specifically for the NPS vegetation and mapping program so that the electronic data entry fields mirror the standard field form. Data entry was facilitated by assigning each plant taxon a unique, standardized code and name based on the PLANTS database developed by Natural Resources Conservation Service in cooperation with the Biota of North America Program (USDA and NRCS 2009). Data was thoroughly proofed after entry to minimize errors.




Figure 3. Locations of all vegetation plots collected at TAPR in 2008.

Plots were assigned to categories based on similarity of vegetation. These categories were assigned names following descriptions in Lauver et al. (1999) and NatureServe Explorer (NatureServe 2006). Where the observed TAPR vegetation did not fit descriptions of natural associations described for Kansas, semi-natural and disturbed associations or alliances described for other parks were considered. In this manner, TAPR vegetation was assigned to one of ten plant associations and alliances.

Once the associations were finalized, a dichotomous key was developed by KBS for use during the Accuracy Assessment (Appendix C). The full NVC hierarchical classification and global descriptions are available in the results section. In addition, the final associations were linked to map classes for use in the photo-interpretation and mapping portions of the project.

In the future, TAPR classification plot data will be used by NatureServe and KBS to update and improve world-wide (i.e., global) descriptions of the NVC plant associations. TAPR specific (i.e., local) descriptions were written based on TAPR plot and AA data. The final TAPR classification contains six NVC vegetation classes and two Park Special vegetation classes.

Digital Imagery and Interpretation

The mapping component was initiated by photo interpretation and digitization of 4-band 2005 IKONOS imagery for the vegetation and land use classes determined through the field visit and expert knowledge of project team members. The heads-up digitization was performed at a display scale of 1:1500 to 1:2000. The digitization, evaluation, and modifications comprised an iterative and collaborative process involving the GIS analysts and the rest of the team. Preliminary maps were checked, corrected, and rechecked for boundary delineations.

Because land management practices, particularly grazing and burning, vegetation phenology, and moisture conditions reveal or mask target map features in the imagery, multiple image sources were used in an attempt to extract the most accurate and comprehensive feature classification possible. There was no one image that captured all vegetation communities and features at their peak differentiability. As noted above, a fall 2005 IKONOS image provided by the NPS served as the general basemap, with additional images used to check mapped features and make adjustments as needed. These image sources included 2003, 2004, 2005, 2006 (3-band, Red, Green, Blue), and 2008 (4-band, Red, Green, Blue, Near Infrared) USDA FSA NAIP, 2002 and 1991 USGS DOQQs, and 1938 panchromatic imagery. Two additional 4-band (Red, Green, Blue, Near Infrared) IKONOS scenes were acquired in June and September of 2008 specifically for this project. These IKONOS image acquisitions targeted specific phenological windows that were generally not covered by other imagery.

The 4-band imagery can be displayed as natural color (Red, Green, Blue) or color infrared imagery. Color infrared is often called false-color because the objects that are normally red appear green, green objects (except vegetation) appear blue, and "infrared" objects appear red. Because healthy green vegetation is a very strong reflector of infrared radiation, and appears bright red in color infrared imagery, it helps tremendously in vegetation mapping efforts. By using color infrared imagery, subtle differences between cool and warm season grasses, wetland vegetation and deciduous trees are apparent and can be accurately delineated.

Geologic strata data layers provided by the Kansas Geological Survey, generalized to shale and limestone, also served as a reference for indentifying zones where thin soiled tallgrass prairie might be present (Sawin, B.S. 2008).

Polygons were assigned map class number and name. The vegetation community polygons and other related and supporting data were then incorporated into a geodatabase format.

Accuracy Assessment

Once the vegetation layer was finalized, the accuracy assessment (AA) was conducted. Typically in mapping exercises both thematic or attribute map accuracy and the positional or polygon line accuracy are considered. In the case of the USGS-NPS National Vegetation Mapping Program, however, the positional accuracy is usually omitted since rarely does vegetation split on discrete edges that can be positively located in the field. The subjectivity involved in this effort plus the high resolution and accuracy of the NAIP and IKONOS basemaps usually allows for the assumption that all products derived from them are well within National Map Accuracy Standards for 1:12,000-scale maps (± 30 feet).

The thematic accuracy of the vegetation map was assessed following the standards provided by the USGS-NPS National Vegetation Mapping Program's Accuracy Assessment Procedures manual (TNC et al. 1994). Assessment included a four step process consisting of a sample design, sample site selection, data collection and data analysis. The design of the AA process followed the five possible scenarios provided in the field manual with stratified random targets placed in each map class based on their respective frequency and abundance (Table 1).

These parameters were loaded into a GIS program along with the vegetation layer. Hawth's Analysis Tools for ArcGIS (Beyer 2004) was used to pick the random target locations and also buffer them 10 meters from any polygon boundary and 50 meters from any other point. Being able to choose minimum distance to polygon boundaries helped to minimize confusion and accounted for the horizontal error typically encountered in common GPS receivers (± 5 m). The resulting target locations were restricted to the authorized boundaries of TAPR due to private land access constraints.

Once the target locations were selected they were downloaded to Garmin or Trimble GPS receivers and investigators walked to the AA points to complete the assessment. During the course of the field work, the estimated position error readings on GPS receivers ranged from 1-7 meters. KBS botanists were provided with draft field maps, map unit definitions, and a key to the associations and alliances (Appendix C). In July 2009, KBS botanists traveled to 132 AA target sites and determined the vegetation association using the field key (Figure 4). At each target they recorded vegetation data on an AA form. They also recorded height and cover of vegetative strata, environmental data, and percent canopy cover of the major species (see AA point form in Appendix D). A rationale for the choice of dominant association was noted when the decision was not clear cut.

Twenty of these points represented springs and seeps, which are not included in the vegetation classification layer. The data recorded on the field forms were subsequently entered into the PLOTS database and reviewed for data entry errors by Kansas Natural Heritage Inventory staff. The results were imported from the database into a GIS layer where they were visually compared in two stages to the vegetation map coverage. The first step was to compare the AA points to the original target locations to check for erroneous points. However, no GPS receiver or location errors were observed.

Table 1. Target number of AA samples per map class based on number of polygons and area.

Scenario	Description	Polygons in class	Area occupied by class	Recommended number of samples in class
Scenario A:	The class is abundant. It covers more than 50 hectares of the total area and consists of at least 30 polygons. In this case, the recommended sample size is 30.	>30	>50 ha	30
Scenario B:	The class is relatively abundant. It covers more than 50 hectares of the total area but consists of fewer than 30 polygons. In this case, the recommended sample size is 20. The rationale for reducing the sample size for this type of class is that sample sites are more difficult to find because of the lower frequency of the class.	<30	>50 ha	20
Scenario C:	The class is relatively rare. It covers less than 50 hectares of the total area but consists of more than 30 polygons. In this case, the recommended sample size is 20. The rationale for reducing the sample size is that the class occupies a small area. At the same time, however, the class consists of a considerable number of distinct polygons that are possibly widely distributed. The number of samples therefore remains relatively high because of the high frequency of the class.	>30	<50 ha	20
Scenario D:	The class is rare. It has more than 5 but fewer than 30 polygons and covers less than 50 hectares of the area. In this case, the recommended number of samples is 5. The rationale for reducing the sample size is that the class consists of small polygons and the frequency of the polygons is low. Specifying more than 5 sample sites will therefore probably result in multiple sample sites within the same (small) polygon. Collecting 5 sample sites will allow an accuracy estimate to be computed, although it will not be very precise.	5-30	<50 ha	5
Scenario E:	The class is very rare. It has fewer than 5 polygons and occupies less than 50 hectares of the total area. In this case, it is recommended that the existence of the class be confirmed by a visit to each sample site. The rationale for the recommendation is that with fewer than 5 sample sites (assuming 1 site per polygon) no estimate of level of confidence can be established for the sample (the existence of the class can only be confirmed through field checking).	<5	<50 ha	Visit all and confirm

Table 2. Summary of the AA statistics used at TAPR.

Statistic	Description
User's Accuracy	The fraction of the accuracy assessment observations in a map class that were found to have the correct vegetation class in the field.
Producer's Accuracy	The fraction of the accuracy assessment observations in a vegetation class in the field that were found to be mapped correctly.
Overall Accuracy	The fraction of accuracy assessment observations within all map classes that were correctly mapped.
Kappa Index	Another measure of overall accuracy, which takes into account the probability that mapped polygons will be correct due to random chance.

The second review step involved comparing the vegetation classification assigned by the field botanists to the vegetation classification assigned to the mapped polygon. If a mismatch was found, the mapped polygon would be corrected.

In the case of TAPR, the AA process was streamlined using methods developed from previous studies at Rocky Mountain National Park (Salas et al. 2004) and Wupatki National Monument (Hansen et al. 2004). All of the statistics and calculations used to analyze these data are described at length in the program manuals (TNC et al. 1994) and are summarized in Table 2. Final assessments for each point were recorded using an error matrix.




Figure 4. Locations of all accuracy assessment points collected at TAPR in 2009.

Results

Vegetation Classification

The final classification for TAPR resulted in eight vegetation classes, six of which had NVC descriptors. The other two classes, restored prairie and weedy vegetation, were considered park specific classifications. The classification results reflect both the moderate amount of diversity of vegetation in the park and a respectably high number of native species. During the sampling efforts a total of 322 species were recorded (Appendix E).

Digital Imagery and Interpretation

For TAPR, 12 map units were developed and directly matched to corresponding plant associations and land-use classes (Table 3). The types included eight vegetation based map units and four land-use classes.

Vegetation Map

Just over 18,800 acres including 10,894 acres in the authorized boundary of TAPR and an additional 7,911 acres in the environs were mapped using 12 map classes (Figure 5). This included four land cover classes and eight vegetation classes. Native Tallgrass Prairie (*Andropogon gerardii* - *Sorghastrum nutans* - *Schizachyrium scoparium* Flint Hills Herbaceous Vegetation) was the most abundant map unit in terms of area, covering 13,758 acres (5,568 hectares) or about 73% of the project area. In terms of frequency, Rocky Mixed Prairie (*Schizachyrium scoparium* - *Bouteloua curtipendula* - *Bouteloua gracilis* Central Plains Herbaceous Vegetation) was most abundant with 286 polygons. Frequencies for each map unit (i.e., number of polygons) and acreage per map unit are listed in Table 3.

The standard minimum mapping unit for NPS vegetation mapping projects is defined as 0.5 hectare. The average area of polygons for this project was 33.0 acres (13.4 hectares).

Accuracy Assessment

The 2009 accuracy assessment effort yielded 112 points that were distributed throughout TAPR; no points were sampled in the environs due to access constraints.

During analysis of the AA points, a GIS point file was created from the AA point coordinates recorded in the field. These were then overlaid on the vegetation map and a comparison of the final AA field call versus the vegetation polygon label was conducted.

Examination of the final error matrix (Appendix A) shows an overall accuracy of 92.0%. Only one map class, cropland, fell below the 80% standard due to its low frequency and low sample size. Areas of confusion occurred between similar vegetation types. Confusion occurred between bur oak forest, successional forest, and wet ravine vegetation, which all had significant overlap in species and were sometimes difficult to distinguish from aerial imagery.

**USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve**

Table 3. Map units identified at TAPR, with their total frequency and acreage.

NVC Identifier	Common Name	Scientific Name / Description	Frequency	Acres	Hectares
Forest and Woodlands					
CEGL002053	Bur Oak Woodland	<i>Quercus macrocarpa</i> / <i>Andropogon gerardii</i> / <i>Hesperostipa spartea</i> Woodland	28	509.8	206.3
CEGL002014	Successional Forest	<i>Fraxinus pennsylvanica</i> / <i>Ulmus</i> spp. / <i>Celtis occidentalis</i> Forest	29	409.0	165.5
Herbaceous Vegetation					
CEGL002201	Native Tallgrass Prairie	<i>Andropogon gerardii</i> - <i>Sorghastrum nutans</i> - <i>Schizachyrium scoparium</i> Flint Hills Herbaceous Vegetation	29	13,758.1	5,567.7
CEGL002246	Rocky Mixed Prairie	<i>Schizachyrium scoparium</i> - <i>Bouteloua curtipendula</i> - <i>Bouteloua gracilis</i> Central Plains Herbaceous Vegetation	286	1,351.6	547.0
CEGL002223	Wet Ravine Vegetation	<i>Spartina pectinata</i> - <i>Eleocharis</i> spp. - <i>Carex</i> spp. Herbaceous Vegetation	21	285.7	115.6
(No assigned code)	Restored Prairie	Planted Semi-natural Restored Tallgrass Prairie, areas that were restored to a tallgrass prairie mix of species	6	97.6	39.5
(No assigned code)	Weedy	Areas of disturbed vegetation, former feedlot	6	319.6	129.3
CEGL005264	Smooth Brome	<i>Bromus inermis</i> - (<i>Pascopyrum smithii</i>) Semi-natural Herbaceous Vegetation	23	559.4	226.4
Land Use/Land Cover					
(No assigned code)	Cropfields	Cultivated fields	42	653.8	264.6
(No assigned code)	Developed Land	Buildings and adjacent lands	24	449.8	182.0
(No assigned code)	Ponds/Water Bodies	Man-made impoundments	73	135.8	55.0
(No assigned code)	Roadways	Highways, county roads, and rights-of-way	2	274.6	111.1
Total Land Use/Land Cover			164	2,073.4	839.1
Total Natural Vegetation			405	16,731.4	6,771.0
Totals			569	18,804.8	7,610.0




Figure 5. Vegetation map of Tallgrass Prairie National Preserve and environs.

Vegetation Associations

(note additional data and explanation can be found at NatureServe, see:
<http://www.natureserve.org/explorer/servlet/NatureServe?init=Ecol>

Mapped Unit Name: *Bur Oak Woodland*

Common Name: Western Tallgrass Bur Oak Woodland

Scientific Name: *Quercus macrocarpa* / *Andropogon gerardii* - *Hesperostipa spartea*
Woodland

Bur Oak / Big Bluestem - Porcupine Grass Woodland

NVC Identifier: CEGL002053



Figure 6. Bur Oak Woodland at Tallgrass Prairie National Preserve, July 2009.

Global Summary: This bur oak woodland community (Figure 6) is associated with the floodplains of rivers and streams in the central-western tallgrass region of the midwestern United States. Stands occur on gentle to steep slopes with silt or loam soils. Soils are well-drained to moderately well-drained, sometimes shallow (0-40 cm), and formed from loess or glacial till. The overstory of this community is open and dominated by *Quercus macrocarpa* (bur oak). *Quercus muehlenbergii* (chinkapin oak) can be a common associate. Shrubs are absent to common, and include *Cornus drummondii* (roughleaf dogwood), *Ceanothus herbaceus* (New Jersey tea), *Corylus americana* (American hazelnut), *Prunus americana* (American plum), *Rhus glabra* (smooth sumac), *Ribes missouriense* (Missouri gooseberry), *Symphoricarpos occidentalis* (western snowberry), and *Zanthoxylum americanum* (common pricklyash). The herbaceous

stratum can be similar to dry prairie. It includes the grasses *Andropogon gerardii* (big bluestem), *Schizachyrium scoparium* (little bluestem), *Sorghastrum nutans* (Indiangrass), *Sporobolus heterolepis* (prairie dropseed), and *Hesperostipa spartea* (porcupinegrass), as well as *Maianthemum stellatum* (starry false lily of the valley), *Monarda fistulosa* (wild bergamot), and *Solidago canadensis* (Canada goldenrod). In the past, periodic fires kept the canopy from closing. Where fire regimes have been disrupted, this community often begins to succeed to other, more closed oak types.

Global Environmental Description: This community occurs near floodplains and on gently sloping to steep upland mesic sites especially within 30 km of the Missouri River but possibly along other rivers. In Nebraska it may have been most abundant in the southeast because conditions are most suitable for tree growth there. The soils on which this community occurs are silt or loam, shallow to deep, with a pH range from 5.6-7.3. The soils of this community are moderately well-drained to well-drained. The parent material is loess or glacial till (Nelson 1987, Lauver et al. 1999, Steinauer and Rolfsmeier 2000).

Global Vegetation Description: The overstory of this community is open and dominated by *Quercus macrocarpa* (bur oak). *Quercus muehlenbergii* (chinkapin oak) can be a common associate. Shrubs are absent to common and include *Cornus drummondii* (roughleaf dogwood), *Ceanothus herbaceus* (New Jersey tea), *Corylus americana* (American hazelnut), *Prunus americana* (American plum), *Rhus glabra* (smooth sumac), *Ribes missouriense* (Missouri gooseberry), *Symphoricarpos occidentalis* (western snowberry), and *Zanthoxylum americanum* (common pricklyash). The herbaceous stratum can be similar to dry prairie. It includes the grasses *Andropogon gerardii* (big bluestem), *Schizachyrium scoparium* (little bluestem), *Sorghastrum nutans* (Indiangrass), *Sporobolus heterolepis* (prairie dropseed), and *Hesperostipa spartea* (porcupinegrass), as well as *Maianthemum stellatum* (starry false lily of the valley), *Monarda fistulosa* (wild bergamot), and *Solidago canadensis* (Canada goldenrod). In the past, periodic fires kept the canopy from closing. Where fire regimes have been disrupted, this community often begins to succeed to other, more closed oak types (Nelson 1987, Lauver et al. 1999, Steinauer and Rolfsmeier 2000).

Most Abundant Species:

Table 4. Overstory Tree Species (DBH>15 cm) within Bur Oak Woodland. (Data collected by Heartland Inventory and Monitoring Program, n=10 plots).

Scientific Name	Common Name	Total Trees
<i>Celtis occidentalis</i>	common hackberry	10
<i>Fraxinus americana</i>	white ash	3
<i>Platanus occidentalis</i>	American sycamore	3
<i>Carya texana</i>	black hickory	2
<i>Fraxinus pennsylvanica</i>	green ash	1
<i>Gymnocladus dioicus</i>	Kentucky coffeetree	1
<i>Populus deltoides</i>	eastern cottonwood	1
<i>Quercus macrocarpa</i>	bur oak	1
<i>Tilia americana</i>	American basswood	1
<i>Ulmus americana</i>	American elm	1
<i>Ulmus rubra</i>	slippery elm	1

Global Conservation Status Rank & Reasons: G2G3. This community has been highly degraded in the mesic sites where it occurred historically. Bur oak woodlands and forests have recently spread upslope into drier areas in the absence of fires. Sites do occur in Missouri in association with loess hill prairies, but are not tracked for conservation purposes because of low quality (M. Leahy pers. comm. 1999).

Mapped Unit Name: *Successional Forest*

Common Name: Central Green Ash - Elm - Hackberry Forest

Scientific Name: *Fraxinus pennsylvanica* - *Ulmus* spp. - *Celtis occidentalis* Forest
Green Ash - Elm species - Common Hackberry Forest

NVC Identifier: C EGL002014



Figure 7. Successional Forest at Tallgrass Prairie National Preserve, July 2009.

Global Summary: This community (Figure 7) is found in the central United States along upper floodplain terraces of rivers and streams and in upland ravine bottoms. Soils are moderately well-drained to poorly drained. Tree canopies are dominated by *Fraxinus pennsylvanica* (green ash), *Celtis occidentalis* (common hackberry), and *Ulmus americana* (American elm). Other tree species that may be present include *Juglans nigra* (black walnut), *Tilia americana* (American basswood), *Acer saccharinum* (silver maple), and *Populus deltoides* (eastern cottonwood). *Ulmus rubra* (slippery elm) can be part of the subcanopy. The shrub layer in the western part of the range includes *Cornus drummondii* (roughleaf dogwood), *Ribes missouriense* (Missouri gooseberry), *Symphoricarpos occidentalis* (western snowberry), and *Zanthoxylum americanum* (common pricklyash), as well as woody vines, such as *Parthenocissus vitacea* (woodbine), *Smilax tamnoides* (bristly greenbrier), *Toxicodendron radicans* (eastern poison-ivy), and *Vitis riparia* (riverbank grape). The herbaceous layer in the western part of its range includes *Elymus virginicus* (Virginia wildrye), *Festuca subverticillata* (nodding fescue), *Galium aparine* (stickywilly), *Geum canadense* (white avens), and *Laportea canadensis* (Canadian woodnettle).

Global Environmental Description: Stands occur along upper floodplain terraces of rivers and streams and in upland ravine bottoms. Soils are moderately well-drained to poorly drained.

Global Vegetation Description: The vegetation has an open to closed tree canopy that is dominated by *Fraxinus pennsylvanica* (green ash), *Celtis occidentalis* (common hackberry), and *Ulmus americana* (American elm). Other tree species that may be present include *Juglans nigra* (black walnut), *Tilia americana* (American basswood), *Acer saccharinum* (silver maple), and *Populus deltoides* (eastern cottonwood). *Ulmus rubra* (slippery elm) can be part of the subcanopy. The shrub layer in the western part of the range includes *Cornus drummondii* (roughleaf dogwood), *Ribes missouriense* (Missouri gooseberry), *Symphoricarpos occidentalis* (western snowberry), and *Zanthoxylum americanum* (common pricklyash), as well as woody vines, such as *Parthenocissus vitacea* (woodbine), *Smilax tamnoides* (bristly greenbrier), *Toxicodendron radicans* (eastern poison-ivy), and *Vitis riparia* (riverbank grape). The herbaceous layer in the western part of the range includes *Elymus virginicus* (Virginia wildrye), *Festuca subverticillata* (nodding fescue), *Galium aparine* (stickywilly), *Geum canadense* (white avens), and *Laportea canadensis* (Canadian woodnettle) (Steinauer and Rolfsmeier 2000).

Most Abundant Species:

Table 5. Overstory Tree Species (DBH>15 cm) within Successional Forest. (Data collected by the Heartland Network breeding bird monitoring (n=4 plots).

Scientific Name	Common Name	Total Trees
<i>Celtis occidentalis</i>	common hackberry	9
<i>Fraxinus americana</i>	white ash	1
<i>Robinia pseudoacacia</i>	black locust	1
<i>Ulmus americana</i>	American elm	1
<i>Ulmus rubra</i>	slippery elm	1

Global Conservation Status Rank & Reasons: G3G5.

Mapped Unit Name: *Native Tallgrass Prairie*

Common Name: Flint Hills Tallgrass Prairie

Scientific Name: *Andropogon gerardii* - *Sorghastrum nutans* - *Schizachyrium scoparium*
Flint Hills Herbaceous Vegetation
Big Bluestem - Yellow Indiangrass - Little Bluestem Flint Hills
Herbaceous Vegetation

NVC Identifier: CEGL002201



Figure 8. Native Tallgrass Prairie at Tallgrass Prairie National Preserve, July 2009.

Global Summary: This tallgrass prairie grassland (Figure 8) is found in the Flint Hills region of the central United States. Stands occur on shallow to deep silt, loam, and clay soils. It can be somewhat poorly drained to somewhat excessively drained. This community has a dense cover of tall grasses with a moderate to high diversity of forbs. Dominant grasses are *Andropogon gerardii* (big bluestem), *Sorghastrum nutans* (Indiangrass), and *Schizachyrium scoparium* (little bluestem). *Bouteloua curtipendula* (sideoats grama), *Panicum virgatum* (switchgrass), and *Sporobolus compositus* (composite dropseed) are common, but less abundant, members of this community. Typical forbs include *Symphotrichum ericoides* (white heath aster), *Helianthus grosseserratus* (sawtooth sunflower), *Lespedeza capitata* (roundhead lespedeza), *Solidago* (goldenrod) spp., and *Viola pedatifida* (prairie violet). Shrubs, such as *Amorpha canescens* (leadplant), and trees are usually infrequent, but can be more common near watercourses.

Global Environmental Description: This community is found on shallow to deep silt, loam, and clay soils. It can be somewhat poorly drained to somewhat excessively drained. The parent material is calcareous clayey shale, limestone, cherty limestone, or interbedded limestone and clayey shale (Lauver et al. 1999).

Global Vegetation Description: This community has a dense cover of tall grasses with a moderate to high diversity of forbs. Dominant grasses are *Andropogon gerardii* (big bluestem), *Sorghastrum nutans* (Indiangrass), and *Schizachyrium scoparium* (little bluestem). *Bouteloua curtipendula* (sideoats grama), *Panicum virgatum* (switchgrass), and *Sporobolus compositus* (composite dropseed) are common, but less abundant, members of this community. Typical forbs include *Symphyotrichum ericoides* (white heath aster), *Helianthus grosseserratus* (sawtooth sunflower), *Lespedeza capitata* (roundhead lespedeza), *Psoralidium tenuiflorum* (slimflower scurfpea), *Solidago* (goldenrod) spp., and *Viola pedatifida* (prairie violet). Shrubs, such as *Amorpha canescens* (leadplant), and trees are usually infrequent, but can be more common near watercourses (Lauver et al. 1999).

Most Abundant Species:

Table 6. Average percent cover of the top twenty most common species in plots within Native Tallgrass Prairie. (Data collected by the Heartland Network in 2008, n=23 plots.)

Scientific Name	Common Name	Average % Cover
<i>Schizachyrium scoparium</i>	little bluestem	23.83
<i>Andropogon gerardii</i>	big bluestem	12.25
<i>Sorghastrum nutans</i>	Indiangrass	9.04
<i>Amphiachyris dracunculoides</i>	prairie broomweed	6.22
<i>Sporobolus compositus</i>	composite dropseed	3.13
<i>Buchloe dactyloides</i>	buffalograss	2.93
<i>Panicum virgatum</i>	switchgrass	2.57
<i>Bouteloua curtipendula</i>	sideoats grama	2.09
<i>Carex spp.</i>	sedge	1.83
<i>Ambrosia psilostachya</i>	Cuman ragweed	1.72
<i>Artemisia ludoviciana</i>	white sagebrush	1.55
<i>Lespedeza violacea</i>	violet lespedeza	0.92
<i>Oxalis spp.</i>	woodsorrel	0.86
<i>Bouteloua hirsuta</i>	hairy grama	0.86
<i>Oxalis violacea</i>	violet woodsorrel	0.82
<i>Bouteloua gracilis</i>	blue grama	0.75
<i>Callirhoe alcaeoides</i>	light poppymallow	0.73
<i>Dichanthelium spp.</i>	rosette grass	0.67
<i>Panicum capillare</i>	witchgrass	0.62

Global Conservation Status Rank & Reasons: G4.

Mapped Unit Name: *Rocky Mixed Prairie*

Common Name: Central Great Plains Little Bluestem Prairie

Scientific Name: *Schizachyrium scoparium* - *Bouteloua curtipendula* - *Bouteloua gracilis*
Central Plains Herbaceous Vegetation
Little Bluestem - Sideoats Grama - Blue Grama Central Plains Herbaceous
Vegetation

NVC Identifier: CEGL 002246



Figure 9. Rocky Mixed Prairie at Tallgrass Prairie National Preserve, July 2009.

Global Summary: This little bluestem - sideoats grama grassland community (Figure 9) is found in the south-central Great Plains of the United States. Stands occur on level to moderately sloping uplands, but are more likely to be on steep ravine slopes. The loam, clay loam, silty loam, or silty soils are usually formed over limestone. They are shallow to moderately deep, well-drained, and usually contain a substantial amount of rock fragments. The vegetation often forms two layers, a shorter layer of grasses and a taller layer of mixed grasses and forbs. Cover is moderately dense to dense in most stands. The vegetation is characteristically dominated by three species, *Schizachyrium scoparium* (little bluestem), *Bouteloua curtipendula* (sideoats grama), and *Bouteloua gracilis* (blue grama). The first two are mid grasses and the latter is a short grass. *Schizachyrium scoparium* (little bluestem) is often the tallest dominant grass, reaching 0.5-0.8 m in Oklahoma. *Andropogon gerardii* (big bluestem), *Sporobolus cryptandrus* (sand dropseed), and *Sorghastrum nutans* (Indiangrass) are present, especially on lower slopes. The short grasses *Buchloe dactyloides* (buffalograss) and *Bouteloua hirsuta* (hairy grama) grow on upper slopes and level ground. Forbs include *Ambrosia psilostachya* (Cuman ragweed),

Dalea enneandra (nineanther prairie clover), *Echinacea angustifolia* (blacksamson echinacea), *Liatris punctata* (dotted blazing star), *Calylophus serrulatus* (yellow sundrops), and *Psoralidium tenuiflorum* (slimflower scurfpea).

Global Environmental Description: This community is primarily found on level to moderately sloping uplands, but is more likely to be on steep ravine slopes in western Kansas (Kuchler 1974). The loam, clay loam, silty loam, or silty soils usually formed over limestone. They are shallow to moderately deep, well-drained, and usually contain a substantial amount of rock fragments (Heitschmidt et al. 1970, Johnston 1987).

Global Vegetation Description: The vegetation in this community often forms two layers, a shorter layer of grasses and a taller layer of mixed grasses and forbs (Kuchler 1974). Cover is moderately dense to dense in most stands (Weaver and Albertson 1956). The vegetation is characteristically dominated by three species, *Schizachyrium scoparium* (little bluestem), *Bouteloua curtipendula* (sideoats grama), and *Bouteloua gracilis* (blue grama). The first two are mid grasses and the latter is a short grass. *Schizachyrium scoparium* (little bluestem) is often the tallest dominant grass, reaching 0.5-0.8 m in Oklahoma (Bruner 1931). *Andropogon gerardii* (big bluestem), *Sporobolus cryptandrus* (sand dropseed), and *Sorghastrum nutans* (Indiangrass) are present, especially on lower slopes. The short grasses *Buchloe dactyloides* (buffalograss) and *Bouteloua hirsuta* (hairy grama) grow on upper slopes and level ground. Forbs include *Ambrosia psilostachya* (Cuman ragweed), *Dalea enneandra* (nineanther prairie clover), *Echinacea angustifolia* (blacksamson echinacea), *Liatris punctata* (dotted blazing star), *Calylophus serrulatus* (yellow sundrops), and *Psoralidium tenuiflorum* (slimflower scurfpea).

Most Abundant Species:

Table 7. Average percent cover of the top twenty most common species in plots within Rocky Mixed Prairie. (Data collected by KBS in 2008, n=10 plots.)

Scientific Name	Common Name	Average % Cover
<i>Andropogon gerardii</i>	big bluestem	40.50
<i>Bouteloua hirsuta</i>	hairy grama	5.90
<i>Bouteloua curtipendula</i>	sideoats grama	4.21
<i>Amorpha canescens</i>	leadplant	3.40
<i>Schizachyrium scoparium</i>	little bluestem	2.41
<i>Panicum virgatum</i>	switchgrass	2.20
<i>Dalea purpurea</i>	purple prairie clover	2.14
<i>Rhus glabra</i>	smooth sumac	1.80
<i>Psoralidium tenuiflorum</i>	slimflower scurfpea	1.71
<i>Sorghastrum nutans</i>	Indiangrass	1.61
<i>Vernonia baldwinii</i>	Baldwin's ironweed	1.32
<i>Ambrosia psilostachya</i>	Cuman ragweed	1.03
<i>Euthamia gymnospermoides</i>	Texas goldentop	1.03
<i>Artemisia ludoviciana</i>	white sagebrush	0.32
<i>Sporobolus compositus</i>	composite dropseed	0.32
<i>Tragia betonicifolia</i>	betonyleaf noseburn	0.32
<i>Baptisia australis</i>	blue wild indigo	0.20
<i>Monarda fistulosa</i>	wild bergamot	0.20
<i>Croton monanthogynus</i>	prairie tea	0.14

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

Scientific Name	Common Name	Average % Cover
<i>Hymenopappus scabiosaeus</i>	Carolina woollywhite	0.13

Global Rank & Reasons: G2G4. The total number of occurrences is unknown. Seven have been documented in Kansas, where the community is ranked S2(?). Although no other occurrences have been documented, the community is also reported in Oklahoma (S?), where it may be more common.

Mapped Unit Name: *Wet Ravine Vegetation*

Common Name: Southern Great Plains Cordgrass Wet Prairie

Scientific Name: *Spartina pectinata* - *Eleocharis* spp. - *Carex* spp. Herbaceous Vegetation
Prairie Cordgrass - Spikerush species - Sedge species Herbaceous
Vegetation

NVC Identifier: CEGL002223



Figure 10. Wet Ravine at Tallgrass Prairie National Preserve, July 2009.

Global Summary: This wet grassland community (Figure 10) is found in the southern Great Plains on deep, poorly drained soils on level to nearly level sites near lakes, seeps, or alluvial lowlands. The soils are usually inundated for short periods during the year but may be saturated for much of the growing season. In northeastern, central, and western Oklahoma (i.e., excluding the Coastal Plain and the Oklahoma panhandle), this association occurs in floodplains, backswamps, and lake margins. This community is characterized by tall, dense graminoids with moderate forb diversity and few woody species. The dominant species, *Spartina pectinata* (prairie cordgrass), can form near monocultures in some locations. Common species include *Carex annectens* (yellowfruit sedge), *Carex blanda* (eastern woodland sedge), *Eleocharis* (spikerush) spp., *Juncus interior* (inland rush), *Juncus torreyi* (Torrey's rush), *Panicum virgatum* (switchgrass), *Rumex altissimus* (pale dock), and *Verbena hastata* (swamp verbena). Other characteristic species in Oklahoma include *Ammannia coccinea* (valley redstem), *Paspalum laeve* (field paspalum), *Pluchea odorata* (sweetscent), and *Vernonia baldwinii* (Baldwin's ironweed), and in Kansas include *Asclepias incarnata* (swamp milkweed), *Symphytichum*

lanceolatum (white panicle aster), *Baptisia alba* var. *macrophylla* (largeleaf wild indigo), *Helianthus grosseserratus* (sawtooth sunflower), and *Scirpus atrovirens* (green bulrush).

Global Environmental Description: This community is found on deep, poorly drained soils on level to nearly level sites near lakes, seeps, or alluvial lowlands (Kuchler 1974, Johnson and Knapp 1995). The soils are usually inundated for short periods during the year, but may be saturated for much of the growing season.

Global Vegetation Description: This community is characterized by tall, dense graminoids with moderate forb diversity and few woody species. The dominant species, *Spartina pectinata* (prairie cordgrass), can form near monocultures in some locations (Johnson and Knapp 1995). Other common species include *Carex annectens* (yellowfruit sedge), *Carex blanda* (eastern woodland sedge), *Eleocharis* (spikerush) spp., *Juncus interior* (inland rush), *Juncus torreyi* (Torrey's rush), *Panicum virgatum* (switchgrass), *Rumex altissimus* (pale dock), and *Verbena hastata* (swamp verbena). Other characteristic species in Oklahoma include *Ammannia coccinea* (valley redstem), *Paspalum laeve* (field paspalum), *Pluchea odorata* (sweetscent), and *Vernonia baldwinii* (Baldwin's ironweed), and in Kansas include *Asclepias incarnata* (swamp milkweed), *Symphotrichum lanceolatum* (white panicle aster), *Baptisia alba* var. *macrophylla* (largeleaf wild indigo), *Helianthus grosseserratus* (sawtooth sunflower), and *Scirpus atrovirens* (green bulrush) (Lauver et al. 1999).

Most Abundant Species:

Table 8. Average percent cover of the top twenty most common species in plots within Wet Ravines. (Data collected by KBS in 2008, n=10 plots; note many areas have woody encroachment)

Scientific Name	Common Name	Average % Cover
<i>Bromus japonicus</i>	field brome	32.31
<i>Amorpha fruticosa</i>	desert false indigo	23.80
<i>Andropogon gerardii</i>	big bluestem	12.90
<i>Vernonia baldwinii</i>	Baldwin's ironweed	8.32
<i>Paspalum pubiflorum</i>	hairyseed paspalum	8.30
<i>Xanthium strumarium</i>	rough cocklebur	7.61
<i>Ambrosia psilostachya</i>	Cuman ragweed	6.02
<i>Sporobolus compositus</i>	composite dropseed	5.31
<i>Panicum virgatum</i>	switchgrass	3.91
<i>Ambrosia trifida</i>	great ragweed	3.01
<i>Muhlenbergia</i> sp.	muhly	2.51
<i>Solidago canadensis</i>	Canada goldenrod	2.21
<i>Sorghastrum nutans</i>	Indiangrass	1.80
<i>Teucrium canadense</i>	Canada germander	1.80
<i>Spartina pectinata</i>	prairie cordgrass	1.60
<i>Glycyrrhiza lepidota</i>	American licorice	1.50
<i>Leersia oryzoides</i>	rice cutgrass	1.50
<i>Symphoricarpos orbiculatus</i>	coralberry	1.41
<i>Tripsacum dactyloides</i>	eastern gamagrass	1.21
<i>Euthamia gymnospermoides</i>	Texas goldentop	1.11

Global Conservation Status Rank & Reasons: G2G4. There are probably more than 20 occurrences rangewide. Six have been documented in Kansas, where the community is ranked SU. Although no other occurrences have been documented, the community is also reported in Oklahoma (S2). It occurs in four ecoregional subsections and has moderately restrictive environmental requirements.

Mapped Unit Name: *Wet Ravine Vegetation - Seeps and Springs*

Common Name: Southern Great Plains Cordgrass Wet Prairie

Scientific Name: *Spartina pectinata* - *Eleocharis* spp. - *Carex* spp. Herbaceous Vegetation
Prairie Cordgrass - Spikerush species - Sedge species Herbaceous
Vegetation

NVC Identifier: CEGL002223

Note: Seeps (Figure 11) and springs were surveyed separately from wet ravines, although they are included within the same community type. Seeps and springs were mapped as point locations, acquired from the Kansas Geological Survey and were not mapped as polygons.



Figure 11. Seep at Tallgrass Prairie National Preserve, July 2009. Darker green in midground is *Eleocharis* spp., tall seedheads are *Scirpus pendulus*, and dark brown seedheads in foreground and background are *Rumex* spp.

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

Table 9. Average percent cover of the top twenty most common species in plots within Seeps and Springs. (Data collected by KBS in 2008, n=10 plots.)

Scientific Name	Common Name	Average % Cover
<i>Eleocharis spp.</i>	spikerush	19.31
<i>Leersia oryzoides</i>	rice cutgrass	14.40
<i>Spartina pectinata</i>	prairie cordgrass	13.00
<i>Scirpus pendulus</i>	rufous bulrush	12.40
<i>Scirpus atrovirens</i>	green bulrush	10.01
<i>Carex spp.</i>	sedge	8.51
<i>Rorippa nasturtium-aquaticum</i>	watercress	8.01
<i>Juncus torreyi</i>	Torrey's rush	7.92
<i>Panicum virgatum</i>	switchgrass	3.11
<i>Ambrosia psilostachya</i>	Cuman ragweed	2.53
<i>Agrimonia parviflora</i>	harvestlice	1.50
<i>Cyperus setigerus</i>	lean flatsedge	1.50
<i>Amorpha fruticosa</i>	desert false indigo	1.20
<i>Glyceria striata</i>	fowl mannagrass	1.11
<i>Ambrosia artemisiifolia</i>	annual ragweed	1.01
<i>Andropogon gerardii</i>	big bluestem	1.01
<i>Carex frankii</i>	Frank's sedge	1.01
<i>Boehmeria cylindrica</i>	smallspike false nettle	1.00
<i>Carex annectens</i>	yellowfruit sedge	0.91
<i>Elymus virginicus</i>	Virginia wildrye	0.81

Mapped Unit Name: Smooth Brome

Common Name: Smooth Brome Semi-natural Grassland

Scientific Name: *Bromus inermis* - (*Pascopyrum smithii*) Semi-natural Herbaceous Vegetation

Smooth Brome - (Western Wheatgrass) Semi-natural Herbaceous Vegetation

NVC Identifier: CEGL005264

Global Summary: This smooth brome grassland type occurs widely throughout the northern Great Plains, in disturbed montane meadows in the Rocky Mountains, on relatively mesic sites in the semi-arid interior western United States, and perhaps more widely in the midwestern U.S. and Canada. Stands can occur in a wide variety of human-disturbed habitats, including highway rights-of-way, jeep trails, etc. The type is also widely planted for revegetating disturbed land, pasture and hay fields, and has escaped into a variety of habitats, including prairie, riparian grasslands, and mesic mountain meadows. The dominant grass is *Bromus inermis* (smooth brome), a naturalized species from Eurasia that forms moderately dense to dense stands that often develop into monocultures. Other weedy species such as *Cirsium arvense* (Canada thistle) and *Poa pratensis* (Kentucky bluegrass) may occur as well, but native species are generally less than 10% cover. Native species may include mixed-grass prairie and montane meadow grasses, such as *Pascopyrum smithii* (western wheatgrass), *Deschampsia caespitosa* (tufted hairgrass), and *Hesperostipa comata* (needle-and-thread), and sparse, scattered mesic shrubs such as *Symphoricarpos* (snowberry) spp., as well as many others. However, the native species are not conspicuous enough to identify the native plant association that could occupy the site, or the stand would be typed as such.

Global Environmental Description: This smooth brome grassland type occurs widely throughout the northern Great Plains, on relatively mesic sites in the semi-arid interior western United States, and perhaps more widely in the midwestern U.S. and Canada. Stands can occur in a wide variety of human-disturbed habitats, including highway rights-of-way, jeep trails, etc. The type is also widely planted for revegetating disturbed land, pasture and hay fields, and has escaped into a variety of habitats, including prairie, riparian grasslands, and mesic mountain meadows. This community is found at all elevational ranges with best examples occurring on mesic alluvial terraces. *Bromus inermis* (smooth brome) grows best on moist, well-drained, finer-textured loam and clay loams, not heavy clays or sand, and does not tolerate prolonged flooding, however, it does persist quite well on well-drained sandy loam derived from granitic parent material. It also occurs in foothills and plains at lower elevations on relatively mesic sites. It occurs on poorly drained sites to rapidly drained sites with fine-textured alluvial soils derived from shale formations found in Utah. This community persists because it is rhizomatous, and once seeded, with enough moisture, will persist, regardless of elevation, soil or landform.

Global Vegetation Description: This association is dominated by medium-tall (0.5-1 m) graminoids. The dominant grass is *Bromus inermis* (smooth brome), a naturalized species from Eurasia that forms moderately dense to dense stands that often develop into monocultures. Other weedy species, such as *Cirsium arvense* (Canada thistle), *Poa pratensis* (Kentucky bluegrass), and other introduced forage species, may occur as well, but native species are generally less than 10% cover. Native species may include mixed-grass prairie and montane meadow grasses, such

as *Juncus balticus* (Baltic rush), *Pascopyrum smithii* (western wheatgrass), *Deschampsia caespitosa* (tufted hairgrass), and *Hesperostipa comata* (needle-and-thread), and sparse scattered mesic shrubs, such as *Artemisia tridentata ssp. wyomingensis* (Wyoming big sagebrush), *Ericameria nauseosa* (rubber rabbitbrush), and *Symphoricarpos* (snowberry) spp., and ruderal forbs, such as *Heterotheca villosa* (hairy false goldenaster), as well as many others. However, the native species are not conspicuous enough to identify the native plant association that could occupy the site, or the stand would be typed as such.

Most Abundant Species:

Table 10. Average percent cover of the top twenty most common species in plots within Smooth Brome Haymeadows. Note that all species other than *Bromus inermis* are under 3% cover. (Data collected by KBS in 2008, n=10 plots.)

Scientific Name	Common Name	Average % Cover
<i>Bromus inermis</i>	smooth brome	96.50
<i>Erigeron annuus</i>	eastern daisy fleabane	2.61
<i>Convolvulus arvensis</i>	field bindweed	2.12
<i>Physalis pumila</i>	dwarf groundcherry	0.52
<i>Bromus japonicus</i>	field brome	0.23
<i>Solanum carolinense</i>	Carolina horsenettle	0.12
<i>Brickellia eupatorioides</i>	false boneset	0.11
<i>Setaria pumila</i>	yellow foxtail	0.11
<i>Conyza canadensis</i>	Canadian horseweed	0.02
<i>Chamaesyce sp.</i>	sandmat	0.01
<i>Oxalis stricta</i>	common yellow oxalis	0.01
<i>Panicum capillare</i>	witchgrass	0.01
<i>Abutilon theophrasti</i>	velvetleaf	0.01
<i>Amaranthus rudis</i>	roughfruit amaranth	0.01
<i>Ambrosia artemisiifolia</i>	annual ragweed	0.01
<i>Asclepias viridis</i>	green antelopehorn	0.01
<i>Chamaesyce nutans</i>	eyebane	0.01
<i>Gleditsia triacanthos</i>	honeylocust	0.01
<i>Mirabilis nyctaginea</i>	heartleaf four o'clock	0.01
<i>Rumex altissimus</i>	pale dock	0.01

Global Conservation Status Rank & Reasons: GNA (invasive) (17-Jun-1999). This is a naturalized type from Europe and Asia, widely planted for cover, pasture, and hay, and has escaped into a variety of habitats.

Common Name: *Restored Prairie*
Scientific Name: Planted Semi-natural Restored Tallgrass Prairie, areas that were restored to a tallgrass prairie mix of species
NVC Identifier: N/A



Figure 12. Restored Prairie at Tallgrass Prairie National Preserve, July 2008.

Global Summary: This community (Figure 12) has been defined for Tallgrass Prairie National Preserve. At TAPR, managers are attempting to restore former brome fields and weedy areas to tallgrass prairie. Natural vegetation of this community is found throughout the northern tallgrass prairie region of the United States and Canada.

Environmental Description: The fields that have been re-planted with native grasses at Tallgrass Prairie National Preserve were once plowed, and have silt-loam soils.

Vegetation Description: This is a grassland community with dense vegetation dominated by tall grasses 1-2 m tall. The abundance of forbs has not reached the abundance found in native Flint Hills Tallgrass Prairie. *Sorghastrum nutans* (Indiangrass) and *Bromus japonicus* (field brome) are the most abundant grasses in this community. *Conyza canadensis* (Canadian horseweed) and *Ambrosia artemisiifolia* (annual ragweed) are common forbs.

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

Most Abundant Species:

Table 11. Average percent cover of the top twenty most common species in plots within Restored Prairies. (Data collected by KBS in 2008, n=19 plots.)

Scientific Name	Common Name	Average % Cover
<i>Sorghastrum nutans</i>	Indiangrass	21.30
<i>Bromus japonicus</i>	field brome	19.52
<i>Panicum virgatum</i>	switchgrass	15.71
<i>Conyza canadensis</i>	Canadian horseweed	15.02
<i>Andropogon gerardii</i>	big bluestem	6.91
<i>Tripsacum dactyloides</i>	eastern gamagrass	6.11
<i>Ambrosia artemisiifolia</i>	annual ragweed	5.00
<i>Abutilon theophrasti</i>	velvetleaf	4.61
<i>Mollugo verticillata</i>	green carpetweed	4.51
<i>Chamaesyce maculata</i>	spotted sandmat	2.70
<i>Setaria faberi</i>	Japanese bristlegrass	2.60
<i>Setaria viridis</i>	green bristlegrass	1.71
<i>Ambrosia trifida</i>	great ragweed	1.70
<i>Erigeron annuus</i>	eastern daisy fleabane	1.61
<i>Bouteloua curtipendula</i>	sideoats grama	1.41
<i>Lactuca serriola</i>	prickly lettuce	1.32
<i>Lactuca saligna</i>	willowleaf lettuce	1.21
<i>Solanum carolinense</i>	Carolina horsenettle	1.21
<i>Amaranthus palmeri</i>	carelessweed	1.20
<i>Schizachyrium scoparium</i>	little bluestem	1.01

Common Name: Weedy
Scientific Name: Areas of disturbed vegetation, former feedlot
NVC Identifier: N/A



Figure 13. Weedy area at Tallgrass Prairie National Preserve, July 2008.

Global Summary and Environmental Description: This community (Figure 13) has been defined for Tallgrass Prairie National Preserve. At TAPR, these are fields that were formerly plowed or used as feedlots, and have not yet been restored to Flint Hills Tallgrass Prairie.

Vegetation Description: This is a grassland community with dense vegetation dominated by tall grasses 1-2 m tall. Exotic grasses, *Bromus inermis* (smooth brome) and *Bromus japonicus* (field brome), are the most abundant grasses in this community. *Convolvulus arvensis* (field bindweed) and *Erigeron annuus* (eastern daisy fleabane) are common forbs.

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

Most Abundant Species:

Table 12. Average percent cover of the top nineteen most common species in plots within Weedy Areas. (Data collected by KBS in 2008, n=4 plots.)

Scientific Name	Common Name	Average % Cover
<i>Bromus inermis</i>	smooth brome	32.50
<i>Bromus japonicus</i>	field brome	31.26
<i>Convolvulus arvensis</i>	field bindweed	20.75
<i>Ambrosia trifida</i>	great ragweed	15.00
<i>Conyza canadensis</i>	Canadian horseweed	7.75
<i>Erigeron annuus</i>	eastern daisy fleabane	6.25
<i>Kochia scoparia</i>	burningbush	6.25
<i>Rumex crispus</i>	curly dock	3.25
<i>Mirabilis nyctaginea</i>	heartleaf four o'clock	2.00
<i>Sorghum halepense</i>	Johnsongrass	2.00
<i>Cucurbita foetidissima</i>	Missouri gourd	1.75
<i>Phytolacca americana</i>	American pokeweed	1.28
<i>Torilis arvensis</i>	spreading hedgeparsley	1.26
<i>Acalypha ostryifolia</i>	pineland threeseed mercury	1.25
<i>Lactuca serriola</i>	prickly lettuce	1.25
<i>Cirsium altissimum</i>	tall thistle	0.75
<i>Helianthus annuus</i>	common sunflower	0.51
<i>Gleditsia triacanthos</i>	honeylocust	0.50
<i>Physalis longifolia</i>	longleaf groundcherry	0.03

Discussion

Tallgrass Prairie National Preserve combines a unique mix of historically important structures and a significant example of native tallgrass prairie.

Field Survey

The vegetation data presented in this project should be used as a baseline to build upon. New survey work in a timely manner would greatly improve both the classification and mapping efforts. Also, accessing neighboring private lands would allow new plot samples to be obtained, increasing the confidence in these types, thereby strengthening the classification.

NVC Classification

In addition to providing a highly accurate vegetation map of the park and environs, we were able to particularly focus on mapping rocky mixed prairie, a sub-community of native tallgrass prairie that is present on limestone outcrops throughout the park.

Digital Imagery and Interpretation

Multiple sources of imagery were used to digitize the vegetation map, which allowed very thorough examination of subtle vegetation characteristics and photo signatures (e.g., shadows of canopy trees). Analyzing imagery taken over multiple seasons, multiple years, and with multiple color band displays allowed us to map boundaries in fine detail and with high confidence.

Accuracy Assessment

We were able to obtain a 92.0% accuracy for map classes, and a kappa adjustment for chance agreements results in an overall accuracy of 90.6%. Our overall accuracy assessment is well above the 80% required by VMP (taking into account the 90% confidence interval). Individual accuracies also met the 80% requirement, with two exceptions:

(1) Cropland: Users' accuracy for this vegetation class is 67%, with a 90% confidence interval of 5%-128%. Of the three accuracy assessment sites mapped as cropland, one was determined to be smooth brome on the ground. Producers' accuracy is 100%, with a 90% confidence interval of 75%-125%.

(2) Central Green Ash - Elm - Hackberry Forest: Producers' accuracy for this vegetation class is 75%, with a 90% confidence interval of 44%-106%. Of the eight accuracy assessment sites classified as this category on the ground, one was mapped as wet ravine and one was mapped as bur oak forest. These three categories were sometimes difficult to distinguish from overhead imagery. Users' accuracy is 86%, with a 90% confidence interval of 57%-115%.

Future Recommendations

In summary, this project represents the best efforts put forth by a multi-disciplined team over a relatively short period in time. In order to create the best possible "long-term" vegetation classification for TAPR and the most accurate and detailed GIS layer, this project should be viewed as a place to start rather than an end product. Present and future NPS staff are encouraged to scrutinize this project, building from its strengths and bolstering its weaknesses. By keeping in mind that this project was only a snapshot in time, future efforts can help complete our understanding of the vegetation in and around TAPR and how it changes. It is the hope of the producers that the

products presented here will help focus and direct future efforts. The following recommendations are summarized below.

1. The diversity of plant species and dynamic nature of the park with respect to the agricultural aspect warrants periodic **field surveys** by experienced ecologists. The inaccessibility of private lands in the environs should be addressed by seeking permission to sample and verify the vegetation. In this way new plant associations could be discovered and existing types could be updated.
2. Remote sensing does not replace on-the-ground knowledge provided by GPS-linked plots, observations and ground verification. Time and funding limitations curtailed the amount of map **ground-truthing** performed. As opportunities arise, maps should be examined in the field by experienced crews. GPS receiver data and other GIS layers should be used to improve and update the spatial data. This map product should not be viewed as static but should be updated with more current and accurate information.
3. For monitoring purposes, **change over time** could be addressed by similar remote sensing projects. New aerial photos or NAIP imagery could be used in regular intervals to document change. Specifically, new imagery could be used to create up-to-date vegetation layers that could be used to compare changes in both individual vegetation stands and across the entire park.
4. In the future, resource management personnel could link the habitat for **species of concern** to specific associations and map units. These map units could then be used to help locate potential sites of endangered or threatened species in the field or identify areas for non-native plant removal or treatment.

Research Opportunities

Having an accurate and current vegetation classification and map presents many new and exciting research opportunities. These include expanding or linking the GIS layer to derive other information such as fire models, habitat monitoring locations, guides for rare plant surveys, and inventorying areas that likely contain exotic or invasive species. The map could be enhanced by overlaying other existing GIS layers such as geology, hydrology, elevation, and soils. In this manner complex interactions between these layers could be examined to yield important information about growth rates, regeneration after disturbance, biomass distribution, and stream morphology. Finally, through innovative analyses the vegetation layer could possibly be used as a springboard for other ecological studies such as examining how the vegetation interacts with soil chemistry, pollution, archeological sites, weather patterns, etc.

Literature Cited

- Beyer, H. L. 2004. Hawth's Analysis Tools for ArcGIS. Available at <http://www.spataleecology.com/htools>.
- Bruner, W. E. 1931. The vegetation of Oklahoma. Ecological Monographs 1:99-188.
- Daubenmire, R. 1959. A canopy-coverage method of vegetational analysis. Northwest Science 33(1): 42-65.
- Federal Geographic Data Committee (FGDC). 1997. Vegetation classification standard, FGDC-STD-005-1997. Available at: <http://www.fgdc.gov/standards/projects/FGDC-standards-projects/vegetation/vegclass.pdf>
- Federal Geographic Data Committee (FGDC). 1998a. Content standard for digital geospatial metadata, FGDC-STD-001-1998. Available at <http://www.fgdc.gov/metadata/geospatial-metadata-standards>
- Federal Geographic Data Committee (FGDC). 1998b. Spatial data transfer standard, FGDC-STD-002 (modified version ANSI NCITS 20:1998). Available at <http://www.fgdc.gov/standards/projects/FGDC-standards-projects/SDTS/>
- Grossman D. H., D. Faber-Langendoen, A. S. Weakley, M. Anderson, P. Bourgeron, R. Crawford, K. Goodin, S. Landaal, K. Metzler, K. D. Patterson, M. Pyne, M. Reid, and L. Sneddon. 1998. International classification of ecological communities: terrestrial vegetation of the United States. Volume I, The National Vegetation Classification System: development, status, and applications. The Nature Conservancy: Arlington, VA. Available at <http://www.natureserve.org/publications/library.jsp#nspubs>
- Grossman, D. H., K. L. Goodin, X. Li, D. Faber-Langendoen, M. Anderson, and R. Vaughan. 1994. Establishing standards for field methods and mapping procedures. Prepared for the USGS-NPS Vegetation Mapping Program by The Nature Conservancy, Arlington VA, and Environmental Science Research Institute, Redlands, CA.
- Hansen, M., J. Coles, K. Thomas, D. Cogan, M. Ried, J. VonLoh, and K. Schulz. 2004. USGS-NPS National Vegetation Mapping Program: Wupatki National Monument, Arizona; Vegetation Classification and Distribution. Final Report. U.S. Geological Survey Southwest Biological Science Center. Flagstaff, AZ.
- Heitschmidt, R. K., G. K. Hulett, and G. W. Tomanek. 1970. Vegetational map and community structure of a west central Kansas prairie. Southwestern Naturalist 14(3):337-350.
- James, K. M., M. D. DeBacker, G. A. Rowell, J. L. Haack and L. W. Morrison. 2009. [Vegetation community monitoring protocol for the Heartland Inventory and Monitoring Network. Natural Resource Report NPS/HTLN/NRR — 2009/141](#). National Park Service, Fort Collins, Colorado.
- Johnson, S. R., and A. K. Knapp. 1995. The influence of fire on *Spartina pectinata* wetland communities in a northeastern Kansas tallgrass prairie. Canadian Journal of Botany 73:84-90.

- Johnston, B. C. 1987. Plant associations of Region Two: Potential plant communities of Wyoming, South Dakota, Nebraska, Colorado, and Kansas. R2-ECOL-87-2. USDA Forest Service, Rocky Mountain Region. Lakewood, CO. 429 pp.
- Kuchler, A. W. 1974. A new vegetation map of Kansas. *Ecology* 55:586-604 (with map supplement).
- Lauver, C. L., K. Kindscher, D. Faber-Langendoen, and R. Schneider. 1999. A classification of the natural vegetation of Kansas. *The Southwestern Naturalist* 44:421-443.
- Leahy, M. Personal communication. 1999. Missouri Natural Heritage Database, Missouri Department of Conservation, Jefferson City.
- McGregor, R. L., and T. M. Barkley, eds. 1986. *Flora of the Great Plains*. University Press of Kansas, Lawrence, KS.
- NatureServe. 2006. NatureServe Explorer: An online encyclopedia of life [web application]. Version 4.7. NatureServe, Arlington, VA. Available at <http://www.natureserve.org/explorer>.
- Nelson, P. W. 1987 [1985]. *The terrestrial natural communities of Missouri*. Missouri Natural Areas Committee, Jefferson City. 197 pp.
- Peitz, D.G., G.A. Rowell, J.L. Haack, K.M. James, L.W. Morrison, and M.D. DeBacker. 2008. [Breeding Bird Monitoring Protocol for the Heartland Network Inventory and Monitoring Program](#). Natural Resource Report NPS/HTLN/NRR- 2008/044. National Park Service, Fort Collins, Colorado.
- Salas, D., J. Stevens, and K. Schulz. 2004. USGS-NPS National Vegetation Mapping Program: Rocky Mountain National Park. Final Report. U.S. Bureau of Reclamation Remote Sensing and GIS Group Technical Memorandum 8260-05-02. Denver, Colorado.
- Sawin, B.S. 2008. Surficial geology of the Tallgrass Prairie National Preserve, Chase County, Kansas: Kansas Geological Survey, Map M-119A, scale 1:12,000.
- Sawin, R. and R. Buchanan. 2000. Water Quality of Selected Springs--Tallgrass Prairie National Preserve, Chase County, Kansas. Kansas Geological Survey, Open-file Report 2000-01.
- Steinauer, G., and S. Rolfsmeier. 2000. Terrestrial natural communities of Nebraska. Unpublished report of the Nebraska Game and Parks Commission. Lincoln, NE. 143 pp.
- The Nature Conservancy (TNC). 1996. *Methodology for Assessing the Utility of Existing Data for Vegetation Mapping*. Arlington, VA.
- The Nature Conservancy (TNC). 1997. *PLOTS Database System, Version 1.1*. Arlington, VA.

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

- The Nature Conservancy (TNC) and Environmental Systems Research Institute (ESRI). 1994a. NBS/NPS Vegetation Mapping Program: Standardized National Vegetation Classification System. Prepared for the U.S. Department of the Interior, National Biological Survey and National Park Service. Washington, D.C.
- The Nature Conservancy (TNC) and Environmental Systems Research Institute (ESRI). 1994b. NBS/NPS Vegetation Mapping Program: Field Methods for Vegetation Mapping. Prepared for the U.S. Department of the Interior, National Biological Survey and National Park Service. Washington, D.C.
- The Nature Conservancy (TNC), Environmental Systems Research Institute, and National Center of Geographic Information and Analysis. 1994. NBS/NPS Vegetation Mapping Program: Accuracy Assessment Procedures. Prepared for the U.S. Department of the Interior, National Biological Survey and National Park Service. Washington, D.C.
- U.S. Geological Survey. 1999. Map accuracy standards. Fact sheet FS-171-99 (November 1999). Web address: <http://egsc.usgs.gov/isb/pubs/factsheets/fs17199.html>.
- USDA and NRCS. 2009. The PLANTS Database (<http://plants.usda.gov>, 19 November 2009). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
- Weaver, J. E., and F. W. Albertson. 1956. Grasslands of the Great Plains: Their nature and use. Johnsen Publishing Co., Lincoln, NE. 395 pp.
- Wolfenbarger and Nimz, 1996. Spring Hill Ranch, Chase County, Kansas, National Historic Landmark Nomination (March 5, 1996)

Appendix A: Contingency Table for Vegetation Mapping at TAPR

Map Units	Tallgrass Prairie	Rocky Mixed Prairie	Wet Ravine	Success. Forest	Weedy	Bur Oak	Smooth Brome	Cropland	Restored Prairie	Totals	Commission Accuracy	90% Conf. Interval	
												-	+
Tallgrass Prairie	19	1			1					21	90%	78%	103%
Rocky Mixed Prairie	1	25								26	96%	88%	104%
Wet Ravine			14	1		1				16	88%	71%	104%
Success. Forest				6		1				7	86%	57%	115%
Weedy					8		1			9	89%	66%	112%
Bur Oak				1		12				13	92%	76%	108%
Smooth Brome							12			12	100%	96%	104%
Cropland							1	2		3	67%	5%	128%
Restored Prairie									5	5	100%	90%	110%
Totals	20	26	14	8	9	14	14	2	5				
Producer's Error	Omission Accuracy	95%	96%	100%	75%	89%	86%	86%	100%	100%	103 Total Correct Points		
	90% Conf. -	84%	88%	96%	44%	66%	67%	67%	75%	90%	112 Total Points		
	Level +	106%	104%	104%	106%	112%	105%	105%	125%	110%			
Overall Total Accuracy = 92.0% Overall Kappa Index = 90.6% Overall 90% Upper and Lower Confidence Interval = 87.3% and 96.6%													

Instructions on Using the Accuracy Assessment Contingency Table:

The contingency table or error matrix found above presents an array of numbers set out in rows and columns corresponding to a particular vegetation map unit relative to the actual vegetation type as verified on the ground. The column headings represent the vegetation classification as determined in the field and the row headings represent the vegetation classification taken from the vegetation map. The highlighted diagonal indicates the number of points assessed in the field that agree with the map label. Conversely, the inaccuracies of each map unit are described as both errors of inclusion (user's or commission errors) and errors of exclusion (producer's or omission errors). By reading across this table (i.e., rows) one can calculate the percent error of commission, or

USGS-NPS Vegetation Mapping Program
Tallgrass Prairie National Preserve

how many polygons for each map unit were incorrectly labeled when compared to the field data. By reading down the table (i.e., columns) one can calculate the percent error of omission, or how many polygons for that type were left off the map. Numbers “on the diagonal” tell the user how well the map unit was interpreted and how confident they can be in using it. Numbers “off the diagonal” yield important information about the deficiencies of the map including which types were: 1) over- mapped - commission errors on the right or 2) under-mapped - omission errors on the bottom.

Appendix B: Example of a Plot Sampling Form

IDENTIFIERS/LOCATORS

Plot Code _____	
Provisional Community Name _____	
State ____ Site Name _____ Local Site Name _____	
Quad Name _____	
GPS file name _____ Field UTM X _____ m E Field UTM Y _____ m N	
Datum _____ Error +/- _____ m	
<i>please do not complete the following information when in the field</i>	
Corrected UTM X _____ m E Corrected UTM Y _____ m N UTM Zone _____	
Project Name _____ Project Leader _____	
Survey Date _____ Surveyor Lead _____ Surveyors _____	
Taxonomic authority _____	
Directions to Plot _____	
Plot length _____ Plot width _____ Plot area _____	
Plot Photos (y/n) ____ Roll Number _____ Frame Number _____ Plot Permanent (y/n) ____	
Plot representativeness _____	

ENVIRONMENTAL DESCRIPTION

Elevation _____ Slope _____ Aspect _____											
Topographic Position _____											
Cowardin System											
<input type="checkbox"/> Upland <input type="checkbox"/> Riverine <input type="checkbox"/> Palustrine <input type="checkbox"/> Lacustrine	<table border="0"> <tr> <td>Non-Tidal</td> <td>Tidal</td> </tr> <tr> <td><input type="checkbox"/> Permanently Flooded</td> <td>_____</td> </tr> <tr> <td><input type="checkbox"/> Semipermanently Flooded</td> <td><input type="checkbox"/> Saturated</td> </tr> <tr> <td><input type="checkbox"/> Seasonally Flooded</td> <td><input type="checkbox"/> Seasonally Flooded/Saturated</td> </tr> <tr> <td><input type="checkbox"/> Temporarily Flooded</td> <td><input type="checkbox"/> Intermittently Flooded</td> </tr> </table>	Non-Tidal	Tidal	<input type="checkbox"/> Permanently Flooded	_____	<input type="checkbox"/> Semipermanently Flooded	<input type="checkbox"/> Saturated	<input type="checkbox"/> Seasonally Flooded	<input type="checkbox"/> Seasonally Flooded/Saturated	<input type="checkbox"/> Temporarily Flooded	<input type="checkbox"/> Intermittently Flooded
Non-Tidal	Tidal										
<input type="checkbox"/> Permanently Flooded	_____										
<input type="checkbox"/> Semipermanently Flooded	<input type="checkbox"/> Saturated										
<input type="checkbox"/> Seasonally Flooded	<input type="checkbox"/> Seasonally Flooded/Saturated										
<input type="checkbox"/> Temporarily Flooded	<input type="checkbox"/> Intermittently Flooded										

Environmental Comments:	Soil Drainage <input type="checkbox"/> Rapidly drained <input type="checkbox"/> Well drained <input type="checkbox"/> Moderately well drained <input type="checkbox"/> Somewhat poorly drained <input type="checkbox"/> Poorly drained <input type="checkbox"/> Very poorly drained
Soil Comments	Landscape/Landform Comments

VEGETATION DESCRIPTION

Leaf phenology (of dominant stratum)	Leaf Type (of dominant stratum)	Physiognomic class	Cover Scale for Species	Height Scale for Strata
<u>Trees or Shrubs</u>	<u>Broad-leaved</u>	<u>Forest</u>	√ Nearby	01 <0.5 m
<u>Evergreen</u>	<u>Needle-leaved</u>	<u>Woodland</u>	1 0-.01%	02 0.5-1m
<u>Cold-deciduous</u>	<u>Microphyllous</u>	<u>Shrubland</u>	2 .01-1%	03 1-2 m
<u>Drought-deciduous</u>	<u>Graminoid</u>	<u>Dwarf-shrubland</u>	3 1-2%	04 2-5 m
<u>Mixed evergreen - cold-deciduous</u>	<u>Forb</u>	<u>Herbaceous</u>	4 2-5%	05 5-10 m
<u>Mixed evergreen - drought-deciduous</u>	<u>Pteridophyte</u>	<u>Nonvascular</u>	5 5-10%	06 10-15 m
		<u>Sparsely Vegetated</u>	6 10-25%	07 15-20 m
			7 25-50%	08 20-35 m
			8 50-75%	09 35 - 50 m
			9 75-95%	10 >50 m
			10 95-100%	
<u>Herbs</u>				
<u>Annual</u>				
<u>Perennial</u>				

Layer (sublayer-optional)	Height Class	Layer % Cover	Dominant and characteristic Species and Cover Class
T Tree			
T_ _____			
T_ _____			
S Shrub			
S_ _____			
S_ _____			
H Herbaceous			
N Nonvascular			
<i>please see above table for height and cover scales</i>			
Animal Use Evidence			
Natural and Anthropogenic Disturbance Comments			
Other Comments			

Appendix C: Tallgrass Prairie National Preserve Dichotomous Key to Vegetation Associations

1a. Plant community dominated by trees.....	2
1b. Plant community dominated by herbaceous vegetation. If woody plants are present, they are scattered individuals or brush due to lack of recent fire.....	3
2a. Woodland or forest of mixed trees of mixed heights, and mixed ages with no old growth trees.....	
Central Green Ash – Elm – Hackberry Forest CEG00214	
2b. Forest with mix of trees, but includes old-growth bur oaks.....	
Western Tallgrass Bur Oak Woodland CEG002053	
3a. Dominated by planted non-native plants.....	4
3b. Dominated by native grasses and native forbs.....	6
4a. Dominated by non-native grasses and annual forbs (annual forbs >5% cover).....	Weedy
4b. Dominated by non-native grasses or crop; if forbs are present, <5% cover.....	5
5a. Planted brome grass.....	Smooth Brome grass <i>Bromus inermis</i> Semi-natural Herbaceous Alliance
5b. Planted agricultural crops.....	Cropland
6a. Native prairie with forbs.....	7
6b. Replanted or Restored Tallgrass Prairie.....	Planted Semi-natural Restored Tallgrass Prairie
7a. Dominated by plants associated with wet places (<i>Eleocharis</i> , <i>Spartina</i> , <i>Amorpha fruticosa</i>).....	8
7b. Dominated by <i>Andropogon gerardii</i>	9
8a. At a spring or seep.....	Spring or Seep
8b. In a riparian area near a stream.....	Southern Great Plains Cordgrass Wet Prairie CEG002223
9a. Greater than 5% rock cover.....	Central Great Plains Little Bluestem Prairie CEG002246
9b. Less than 5% rock cover.....	Flint Hills Tallgrass Prairie CEG002201

Appendix D: Example of an Accuracy Assessment Sampling Form

NPS Vegetation Mapping: Accuracy Assessment Form

Plot # _____ Park Code: **TAPR** Observers: _____ Date: _____
 UTM X _____ m E UTM Y _____ m N Zone: _____
 Datum: _____ PDOP _____ Elevation _____ Waypoint: _____
 Topography: _____ Slope: _____ Picture no(s): _____

	Stratum	Height (m)	% Cover of Strata	Major Species Present	% Cover of Each Species
T1	Emergent				
T2	Canopy				
T3	Subcanopy (<10 cm DBH)				
S1	Tall Shrub (2-5 m)				
S2	Short Shrub (<2m)				
H	Herbaceous				
A1	Floating Leaved Aquatic				
A2	Submerged Aquatic				
N	Nonvascular				

Comments on indicator species or rare species: _____

Mapped Vegetation Association: _____

Observed Vegetation Association: _____

Comments (note influences on vegetation, difficulties with classification, etc):

Appendix E: Tallgrass Prairie National Preserve Species List

Family	Scientific Name	Common Name
Acanthaceae	<i>Justicia americana</i>	American water-willow
	<i>Ruellia humilis</i>	fringeleaf wild petunia
Aceraceae	<i>Acer negundo</i>	boxelder
Amaranthaceae	<i>Amaranthus palmeri</i>	carelessweed
	<i>Amaranthus tuberculatus</i>	roughfruit amaranth
Anacardiaceae	<i>Rhus aromatica</i>	fragrant sumac
	<i>Rhus glabra</i>	smooth sumac
	<i>Toxicodendron radicans</i>	eastern poison ivy
Apiaceae	<i>Cicuta maculata</i>	spotted water hemlock
	<i>Lomatium foeniculaceum</i>	desert biscuitroot
	<i>Spermolepis inermis</i>	Red River scaleseed
	<i>Torilis arvensis</i>	spreading hedgeparsley
Apocynaceae	<i>Apocynum cannabinum</i>	Indianhemp
Asclepiadaceae	<i>Asclepias sullivantii</i>	prairie milkweed
	<i>Asclepias syriaca</i>	common milkweed
	<i>Asclepias tuberosa</i>	butterfly milkweed
	<i>Asclepias verticillata</i>	whorled milkweed
	<i>Asclepias viridiflora</i>	green comet milkweed
	<i>Asclepias viridis</i>	green antelopehorn
Asteraceae	<i>Achillea millefolium</i>	common yarrow
	<i>Ageratina altissima</i>	white snakeroot
	<i>Ambrosia artemisiifolia</i>	annual ragweed
	<i>Ambrosia psilostachya</i>	Cuman ragweed
	<i>Ambrosia trifida</i>	great ragweed
	<i>Amphiachyris dracunculoides</i>	prairie broomweed
	<i>Antennaria neglecta</i>	field pussytoes
	<i>Arnoglossum plantagineum</i>	groovestem Indian pliantain
	<i>Artemisia ludoviciana</i>	white sagebrush
	<i>Bidens frondosa</i>	devil's beggartick
	<i>Brickellia eupatorioides</i>	false boneset
	<i>Cirsium altissimum</i>	tall thistle
	<i>Cirsium undulatum</i>	wavyleaf thistle
	<i>Conyza canadensis</i>	Canadian horseweed
	<i>Echinacea angustifolia</i>	blacksamson echinacea
	<i>Eclipta prostrata</i>	false daisy
	<i>Erechtites hieraciifolia</i>	American burnweed
	<i>Erigeron annuus</i>	eastern daisy fleabane
	<i>Erigeron philadelphicus</i>	Philadelphia fleabane
	<i>Erigeron strigosus</i>	prairie fleabane
	<i>Eupatorium altissimum</i>	tall thoroughwort
	<i>Euthamia gymnospermoides</i>	Texas goldentop
	<i>Helianthus annuus</i>	common sunflower
	<i>Helianthus grosseserratus</i>	sawtooth sunflower
	<i>Helianthus maximiliani</i>	Maximilian sunflower
	<i>Helianthus pauciflorus</i>	stiff sunflower

Family	Scientific Name	Common Name
	<i>Hieracium longipilum</i>	hairy hawkweed
	<i>Hymenopappus scabiosaeus</i>	Carolina woollywhite
	<i>Iva annua</i>	annual marshelder
	<i>Krigia caespitosa</i>	weedy dwarfdandelion
	<i>Lactuca canadensis</i>	Canada lettuce
	<i>Lactuca saligna</i>	willowleaf lettuce
	<i>Lactuca serriola</i>	prickly lettuce
	<i>Liatris punctata</i>	dotted blazing star
	<i>Nothocalais cuspidata</i>	prairie false dandelion
	<i>Oligoneuron rigidum</i>	stiff goldenrod
	<i>Packera plattensis</i>	prairie groundsel
	<i>Ratibida columnifera</i>	upright prairie coneflower
	<i>Ratibida pinnata</i>	pinnate prairie coneflower
	<i>Rudbeckia hirta</i>	blackeyed Susan
	<i>Silphium integrifolium</i>	wholeleaf rosinweed
	<i>Silphium laciniatum</i>	compassplant
	<i>Solidago canadensis</i>	Canada goldenrod
	<i>Solidago gigantea</i>	giant goldenrod
	<i>Solidago missouriensis</i>	Missouri goldenrod
	<i>Solidago speciosa</i>	showy goldenrod
	<i>Symphyotrichum ericoides</i>	white heath aster
	<i>Symphyotrichum laeve</i>	smooth blue aster
	<i>Symphyotrichum lanceolatum</i>	white panicle aster
	<i>Symphyotrichum oblongifolium</i>	aromatic aster
	<i>Symphyotrichum praealtum</i>	willowleaf aster
	<i>Symphyotrichum sericeum</i>	western silver aster
	<i>Taraxacum officinale</i>	common dandelion
	<i>Verbesina alternifolia</i>	wingstem
	<i>Vernonia baldwinii</i>	Baldwin's ironweed
	<i>Xanthium strumarium</i>	rough cocklebur
Boraginaceae	<i>Lithospermum canescens</i>	hoary puccoon
	<i>Lithospermum incisum</i>	narrowleaf stoneseed
	<i>Myosotis verna</i>	spring forget-me-not
	<i>Onosmodium bejariense</i>	soft-hair marbleseed
Brassicaceae	<i>Alliaria petiolata</i>	garlic mustard
	<i>Camelina microcarpa</i>	littlepod false flax
	<i>Descurainia pinnata</i>	western tansymustard
	<i>Draba brachycarpa</i>	shortpod draba
	<i>Draba cuneifolia</i>	wedgeleaf draba
	<i>Draba reptans</i>	Carolina draba
	<i>Lepidium densiflorum</i>	common pepperweed
	<i>Nasturtium officinale</i>	watercress
	<i>Thlaspi arvense</i>	field pennycress
Cactaceae	<i>Escobaria missouriensis</i>	Missouri foxtail cactus
	<i>Opuntia macrorhiza</i>	twistspine pricklypear
Campanulaceae	<i>Lobelia siphilitica</i>	great blue lobelia
	<i>Triodanis perfoliata</i>	clasping Venus' looking-glass

Family	Scientific Name	Common Name
Caprifoliaceae	<i>Sambucus nigra</i>	European black elderberry
	<i>Symphoricarpos orbiculatus</i>	coralberry
Caryophyllaceae	<i>Cerastium brachypodum</i>	shortstalk chickweed
	<i>Dianthus armeria</i>	Deptford pink
	<i>Silene antirrhina</i>	sleepy silene
	<i>Stellaria media</i>	common chickweed
Chenopodiaceae	<i>Bassia scoparia</i>	burningbush
	<i>Chenopodium album</i>	lambsquarters
Commelinaceae	<i>Tradescantia bracteata</i>	longbract spiderwort
Convolvulaceae	<i>Calystegia sepium</i>	hedge false bindweed
	<i>Convolvulus arvensis</i>	field bindweed
	<i>Evolvulus nuttallianus</i>	shaggy dwarf morning-glory
Cornaceae	<i>Cornus drummondii</i>	roughleaf dogwood
Cucurbitaceae	<i>Cucurbita foetidissima</i>	Missouri gourd
Cupressaceae	<i>Juniperus virginiana</i>	eastern redcedar
Cyperaceae	<i>Carex annectens</i>	yellowfruit sedge
	<i>Carex austrina</i>	southern sedge
	<i>Carex bicknellii</i>	Bicknell's sedge
	<i>Carex brevior</i>	shortbeak sedge
	<i>Carex frankii</i>	Frank's sedge
	<i>Carex gravida</i>	heavy sedge
	<i>Carex molesta</i>	troublesome sedge
	<i>Carex muehlenbergii</i>	Muhlenberg's sedge
	<i>Carex vulpinoidea</i>	fox sedge
	<i>Cyperus acuminatus</i>	tapertip flatsedge
	<i>Cyperus esculentus</i>	yellow nutsedge
	<i>Cyperus lupulinus</i>	Great Plains flatsedge
	<i>Cyperus odoratus</i>	fragrant flatsedge
	<i>Cyperus setigerus</i>	lean flatsedge
	<i>Cyperus strigosus</i>	strawcolored flatsedge
	<i>Eleocharis compressa</i>	flatstem spikerush
	<i>Schoenoplectus tabernaemontani</i>	softstem bulrush
	<i>Scirpus atrovirens</i>	green bulrush
	<i>Scirpus pendulus</i>	rufous bulrush
	Euphorbiaceae	<i>Acalypha ostryifolia</i>
<i>Acalypha rhomboidea</i>		common threeseed mercury
<i>Acalypha virginica</i>		Virginia threeseed mercury
<i>Chamaesyce maculata</i>		spotted sandmat
<i>Chamaesyce nutans</i>		eyebane
<i>Chamaesyce prostrata</i>		prostrate sandmat
<i>Croton capitatus</i>		hogwort
<i>Croton monanthogynus</i>		prairie tea
<i>Euphorbia dentata</i>		toothed spurge
<i>Euphorbia marginata</i>		snow on the mountain
<i>Euphorbia spathulata</i>		warty spurge
<i>Tragia betonicifolia</i>		betonyleaf noseburn
Fabaceae	<i>Amorpha canescens</i>	leadplant

Family	Scientific Name	Common Name
	<i>Amorpha fruticosa</i>	desert false indigo
	<i>Astragalus canadensis</i>	Canadian milkvetch
	<i>Astragalus crassicaarpus</i>	groundplum milkvetch
Fabaceae	<i>Baptisia alba</i>	white wild indigo
	<i>Baptisia australis</i>	blue wild indigo
	<i>Baptisia bracteata</i>	longbract wild indigo
	<i>Cercis canadensis</i>	eastern redbud
	<i>Chamaecrista fasciculata</i>	partridge pea
	<i>Dalea candida</i>	white prairie clover
	<i>Dalea multiflora</i>	roundhead prairie clover
	<i>Dalea purpurea</i>	violet prairie clover
	<i>Desmanthus illinoensis</i>	prairie bundleflower
	<i>Desmodium glutinosum</i>	pointedleaf ticktrefoil
	<i>Desmodium illinoense</i>	Illinois ticktrefoil
	<i>Desmodium sessilifolium</i>	sessileleaf ticktrefoil
	<i>Gleditsia triacanthos</i>	honeylocust
	<i>Glycine max</i>	soybean
	<i>Glycyrrhiza lepidota</i>	American licorice
	<i>Gymnocladus dioicus</i>	Kentucky coffeetree
	<i>Lespedeza capitata</i>	roundhead lespedeza
	<i>Lespedeza violacea</i>	violet lespedeza
	<i>Lotus unifoliolatus</i>	American bird's-foot trefoil
	<i>Medicago lupulina</i>	black medick
	<i>Melilotus officinalis</i>	yellow sweetclover
	<i>Mimosa nuttallii</i>	Nuttall's sensitive-briar
	<i>Pediomelum argophyllum</i>	silverleaf Indian breadroot
	<i>Psoralidium tenuiflorum</i>	slimflower scurfpea
	<i>Robinia pseudoacacia</i>	black locust
	<i>Strophostyles leiosperma</i>	slickseed fuzzybean
Fagaceae	<i>Quercus macrocarpa</i>	bur oak
	<i>Quercus muehlenbergii</i>	chinkapin oak
Fumariaceae	<i>Corydalis micrantha</i>	smallflower fumewort
Gentianaceae	<i>Gentiana puberulenta</i>	downy gentian
Geraniaceae	<i>Geranium carolinianum</i>	Carolina geranium
Hippocastanaceae	<i>Aesculus glabra</i>	Ohio buckeye
Iridaceae	<i>Sisyrinchium campestre</i>	prairie blue-eyed grass
Juglandaceae	<i>Carya texana</i>	black hickory
	<i>Juglans nigra</i>	black walnut
Juncaceae	<i>Juncus dudleyi</i>	Dudley's rush
	<i>Juncus interior</i>	inland rush
	<i>Juncus torreyi</i>	Torrey's rush
Lamiaceae	<i>Hedeoma hispida</i>	rough false pennyroyal
	<i>Lamium amplexicaule</i>	henbit deadnettle
	<i>Lycopus americanus</i>	American water horehound
	<i>Monarda fistulosa</i>	wild bergamot
	<i>Prunella vulgaris</i>	common selfheal
	<i>Salvia azurea</i>	azure blue sage

Family	Scientific Name	Common Name
	<i>Scutellaria parvula</i>	small skullcap
	<i>Teucrium canadense</i>	Canada germander
	<i>Trichostema brachiatum</i>	fluxweed
Lemnaceae	<i>Lemna minor</i>	common duckweed
Liliaceae	<i>Allium spp.</i>	onion
	<i>Erythronium mesochoreum</i>	midland fawnlily
	<i>Nothoscordum bivalve</i>	crowpoison
	<i>Zigadenus nuttallii</i>	Nuttall's deathcamas
Linaceae	<i>Linum sulcatum</i>	grooved flax
Loasaceae	<i>Mentzelia oligosperma</i>	chickenthiel
Lythraceae	<i>Ammannia coccinea</i>	valley redstem
	<i>Lythrum alatum</i>	winged lythrum
Malvaceae	<i>Abutilon theophrasti</i>	velvetleaf
	<i>Callirhoe alcaeoides</i>	light poppymallow
	<i>Sida spinosa</i>	prickly fanpetals
Menispermaceae	<i>Menispermum canadense</i>	moonseed
Molluginaceae	<i>Mollugo verticillata</i>	green carpetweed
Moraceae	<i>Maclura pomifera</i>	osage orange
	<i>Morus alba</i>	white mulberry
	<i>Morus rubra</i>	red mulberry
Nyctaginaceae	<i>Mirabilis albidia</i>	white four o'clock
	<i>Mirabilis nyctaginea</i>	heartleaf four o'clock
Oleaceae	<i>Fraxinus americana</i>	white ash
	<i>Fraxinus pennsylvanica</i>	green ash
Onagraceae	<i>Gaura mollis</i>	velvetweed
	<i>Ludwigia alternifolia</i>	seedbox
	<i>Ludwigia peploides</i>	floating primrose-willow
	<i>Oenothera biennis</i>	common evening-primrose
	<i>Oenothera laciniata</i>	cutleaf evening primrose
	<i>Oenothera macrocarpa</i>	bigfruit evening-primrose
	<i>Oenothera speciosa</i>	pinkladies
Orchidaceae	<i>Spiranthes cernua</i>	nodding lady's tresses
Oxalidaceae	<i>Oxalis stricta</i>	common yellow oxalis
	<i>Oxalis violacea</i>	violet woodsorrel
Phytolaccaceae	<i>Phytolacca americana</i>	American pokeweed
Plantaginaceae	<i>Plantago rhodosperma</i>	redseed plantain
	<i>Plantago rugelii</i>	blackseed plantain
	<i>Plantago virginica</i>	Virginia plantain
Platanaceae	<i>Platanus occidentalis</i>	American sycamore
Poaceae	<i>Alopecurus carolinianus</i>	Carolina foxtail
	<i>Andropogon gerardii</i>	big bluestem
	<i>Aristida oligantha</i>	prairie threeawn
	<i>Bothriochloa laguroides</i>	silver beardgrass
	<i>Bouteloua curtipendula</i>	sideoats grama
	<i>Bouteloua dactyloides</i>	buffalograss
	<i>Bouteloua gracilis</i>	blue grama
	<i>Bouteloua hirsuta</i>	hairy grama

Family	Scientific Name	Common Name
	<i>Bromus arvensis</i>	field brome
	<i>Bromus inermis</i>	smooth brome
	<i>Chloris verticillata</i>	tumble windmill grass
	<i>Dichantherium acuminatum</i>	tapered rosette grass
	<i>Dichantherium oligosanthes</i>	Heller's rosette grass
	<i>Dichantherium wilcoxianum</i>	fall rosette grass
	<i>Digitaria cognata</i>	fall witchgrass
Poaceae	<i>Digitaria sanguinalis</i>	hairy crabgrass
	<i>Echinochloa muricata</i>	rough barnyardgrass
	<i>Elymus canadensis</i>	Canada wildrye
	<i>Elymus virginicus</i>	Virginia wildrye
	<i>Eragrostis spectabilis</i>	purple lovegrass
	<i>Festuca subverticillata</i>	nodding fescue
	<i>Glyceria striata</i>	fowl mannagrass
	<i>Koeleria macrantha</i>	prairie Junegrass
	<i>Leersia oryzoides</i>	rice cutgrass
	<i>Leptochloa panicea</i>	mucronate sprangletop
	<i>Muhlenbergia cuspidata</i>	plains muhly
	<i>Muhlenbergia mexicana</i>	Mexican muhly
	<i>Panicum capillare</i>	witchgrass
	<i>Panicum virgatum</i>	switchgrass
	<i>Paspalum pubiflorum</i>	hairyseed paspalum
	<i>Poa pratensis</i>	Kentucky bluegrass
	<i>Schedonorus phoenix</i>	tall fescue
	<i>Schizachyrium scoparium</i>	little bluestem
	<i>Setaria faberi</i>	Japanese bristlegrass
	<i>Setaria pumila</i>	yellow bristlegrass
	<i>Setaria viridis</i>	green bristlegrass
	<i>Sorghastrum nutans</i>	Indiangrass
	<i>Sorghum halepense</i>	Johnsongrass
	<i>Spartina pectinata</i>	prairie cordgrass
	<i>Sphenopholis obtusata</i>	prairie wedgescale
	<i>Sporobolus compositus</i>	composite dropseed
	<i>Sporobolus heterolepis</i>	prairie dropseed
	<i>Sporobolus neglectus</i>	puffsheath dropseed
	<i>Sporobolus vaginiflorus</i>	poverty dropseed
	<i>Tridens flavus</i>	purpletop tridens
	<i>Tripsacum dactyloides</i>	eastern gamagrass
	<i>Vulpia octoflora</i>	sixweeks fescue
Polygalaceae	<i>Polygala verticillata</i>	whorled milkwort
Polygonaceae	<i>Polygonum amphibium</i>	water knotweed
	<i>Polygonum persicaria</i>	spotted ladythumb
	<i>Rumex altissimus</i>	pale dock
	<i>Rumex crispus</i>	curly dock
Portulacaceae	<i>Portulaca oleracea</i>	little hogweed
Primulaceae	<i>Androsace occidentalis</i>	western rockjasmine
Pteridaceae	<i>Pellaea glabella</i>	smooth cliffbrake

Family	Scientific Name	Common Name
Ranunculaceae	<i>Delphinium carolinianum</i>	Carolina larkspur
	<i>Myosurus minimus</i>	tiny mousetail
Rhamnaceae	<i>Ceanothus americanus</i>	New Jersey tea
	<i>Ceanothus herbaceus</i>	Jersey tea
Rosaceae	<i>Agrimonia parviflora</i>	harvestlice
	<i>Geum canadense</i>	white avens
	<i>Prunus americana</i>	American plum
	<i>Rosa arkansana</i>	prairie rose
Rubiaceae	<i>Galium aparine</i>	stickywilly
Salicaceae	<i>Populus deltoides</i>	eastern cottonwood
	<i>Salix interior</i>	sandbar willow
	<i>Salix nigra</i>	black willow
Santalaceae	<i>Comandra umbellata</i>	bastard toadflax
Scrophulariaceae	<i>Leucospora multifida</i>	narrowleaf paleseed
	<i>Mimulus ringens</i>	Allegheny monkeyflower
	<i>Veronica anagallis-aquatica</i>	water speedwell
	<i>Veronica arvensis</i>	corn speedwell
	<i>Veronica peregrina</i>	neckweed
Solanaceae	<i>Physalis heterophylla</i>	clammy groundcherry
	<i>Physalis longifolia</i>	longleaf groundcherry
	<i>Physalis pumila</i>	dwarf groundcherry
	<i>Physalis virginiana</i>	Virginia groundcherry
	<i>Solanum carolinense</i>	Carolina horsenettle
	<i>Solanum rostratum</i>	buffalobur nightshade
Tiliaceae	<i>Tilia americana</i>	American basswood
Ulmaceae	<i>Celtis occidentalis</i>	common hackberry
	<i>Ulmus americana</i>	American elm
	<i>Ulmus rubra</i>	slippery elm
Urticaceae	<i>Boehmeria cylindrica</i>	smallspike false nettle
	<i>Laportea canadensis</i>	Canadian woodnettle
	<i>Parietaria pensylvanica</i>	Pennsylvania pellitory
	<i>Urtica dioica</i>	stinging nettle
Verbenaceae	<i>Glandularia bipinnatifida</i>	Dakota mock vervain
	<i>Verbena hastata</i>	swamp verbena
	<i>Verbena simplex</i>	narrowleaf vervain
	<i>Verbena stricta</i>	hoary verbena
	<i>Verbena urticifolia</i>	white vervain
Violaceae	<i>Hybanthus verticillatus</i>	babyslippers
	<i>Viola bicolor</i>	field pansy
	<i>Viola nephrophylla</i>	northern bog violet
	<i>Viola pedatifida</i>	prairie violet
Vitaceae	<i>Parthenocissus quinquefolia</i>	Virginia creeper

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 031/107299, April 2011

National Park Service
U.S. Department of the Interior



Natural Resource Program Center
1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525

www.nature.nps.gov

EXPERIENCE YOUR AMERICA™